


NET CONTENTS GALLONS



**CONTAINS
6 LBS TOXAPHENE PER GALLON**

ACTIVE INGREDIENTS

Toxaphene 57.08%
Heavy Aromatic Naphtha 38.92%

INERT INGREDIENTS

4.00%
Total 100.00%

E.P.A. Reg. No. 557-1876
E.P.A. Est. No. 557-FL-1

*Technical Chlorinated Camphene (Chlorine Content 67-69%)

WARNING: Keep Out Of Reach Of Children
See Side Panels For Additional Warnings

DIRECTIONS FOR USE

SMALL GRAINS (Wheat, Barley, Oats, Rye, Rice, Grain Sorghum) For control of armyworms and cutworms, use 1 1/2 quarts of this product suitably diluted with water on one acre of applicable area.

Do not use on treated sorghum forage. Do not apply to sorghum within 30 days of harvest. Do not apply more than one treatment after heads start to form. Do not feed treated forage to dairy animals or animals being finished for slaughter.

CABBAGE For control of cutworms (soil), apply 1 1/2 to 2 pounds of active ingredient per acre, or 1 1/2 to 2 pints of this formulation, suitably diluted by broadcasting, on the soil surface. For control of diamondback moth, imported cabbage worms, and cabbage looper, use 2 to 3 pounds of active ingredient per acre, or 1 1/2 to 2 pints of this formulation, if cabbage worms are present, add 2 1/2 pounds of a dust particle.

Do not apply after heads start to form. Plants treated with this formulation should not be fed to dairy animals or animals being finished for slaughter.

GRAIN AND FORAGE For control of all types and species of armyworms, cutworms, and grasshoppers, apply 1 1/2 to 2 quarts of this product per acre, or 1 1/2 to 2 pints of this formulation, on the soil surface.

For control of cabbage looper, apply at the rate of 2 1/2 to 5 1/2 pints per acre in sufficient water to give good distribution.

Do not make more than a total of 4 applications nor more than 3 applications after thinning. Do not apply at the higher rate within 35 days of harvest. Do not apply at the low rate within 28 days of harvest.

TOMATOES For control of tomato hornworm, tomato fruitworm, Colorado potato beetle, pinworm, armyworm, cutworm, serpentine leaf miner, flea beetle, and russet mite, apply at the rate of 2 1/2 to 6 1/2 pints (2 to 5 pounds of actual toxaphene) per acre, depending on the size of the plants.


Do not apply within 3 days of harvest.

GRASS (Ornamental Lawns and Turf Only) For control of armyworm (true and fall), webworms (including sod and garden webworms), apply at the rate of 3 to 4 pints per acre in sufficient water to give good coverage. For control of grasshoppers, cutworms (including climbing and surface feeding), and grasshoppers, apply at the rate of 2 1/2 pints per acre in sufficient water to give good distribution.

For use only along highway on grass, powerline right of way, picnic areas, home lawns, and other non-agricultural areas or land.

COTTON For control of flea beetles, cotton flea hoppers, thrips, green webworms, and yellow striped armyworms, apply at the rate of 1 1/2 to 2 quarts per acre in sufficient water to give distribution. For control of boll weevils, boll worms, cutworms (including and surface), apply as above at a rate of 1 1/2 to 2 1/2 quarts per acre. For control of fall armyworms, grasshoppers, armyworms, apply at a rate of 1 1/2 to 2 quarts per acre. Do not apply to bolls or open. For control of stink bugs, apply as above at the rate of 1 1/2 quarts per acre.

Do not apply within 30 days of harvest. Apply as needed and repeat at 7 to 14 day intervals until controlled. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not apply to feed forage. Do not use on forage to be used normally or stored.



interstate. Do not store near feed and foodstuffs. Do not treat calves under 3 months of age.

WARNING

May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe spray mist. Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns. Do not use on cucurbits.

ENVIRONMENTAL PRECAUTIONS

This product is toxic to fish, shrimp, crab, birds, and other wildlife. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Destroy all containers when empty by punching holes in them and burying with wastes. Do not burn. Do not reuse containers for other purposes. Rinse equipment and containers, and dispose of wastes by burying in non-crop lands away from water supplies.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

ANTIDOTE

INTERNAL: Give sodium chloride (table salt) in glass of warm water until vomit is clear. Call a doctor immediately.

EXTERNAL: Wash with soap and water.

Manufactured By
Swift Chemical Company
(An E. I. du Pont Company)
Chicago, Illinois 60604