

**ACCEPTED**  
 500 17 18/5  
 U.S. Department of Agriculture,  
 Federal Plant Quarantine Act,  
 Section 17, under  
 P. H. No. 17-18/5

# SWIFT

## COPPER FUNGICIDE SPRAY FOR FLORIDA PLANTS

Plant	To Prevent	When to Apply	Amount to Use
Apples	Apple Blight Apple Scab Apple Rust	5 applications starting when buds are opening and at 4 week intervals.	1.5 lbs per 100 sq ft of leaves or 1.0 lb per 100 sq ft of fruit
Oranges	Anthracnose Scab	5 applications starting when first leaves are opening and at 4 week intervals.	1.5 lbs per 100 sq ft of leaves or 1.0 lb per 100 sq ft of fruit
Limes	Anthracnose Scab	5 applications starting when first leaves are opening and at 4 week intervals.	1.5 lbs per 100 sq ft of leaves or 1.0 lb per 100 sq ft of fruit
Guavas	Anthracnose Scab	5 applications starting when first leaves are opening and at 4 week intervals.	1.5 lbs per 100 sq ft of leaves or 1.0 lb per 100 sq ft of fruit
Avocados	Black Spot	When disease first appears, spray at 4 week intervals and at 4 week intervals.	1.5 lbs per 100 sq ft of leaves or 1.0 lb per 100 sq ft of fruit
Peaches	Black Spot	When disease first appears, spray at 4 week intervals and at 4 week intervals.	1.5 lbs per 100 sq ft of leaves or 1.0 lb per 100 sq ft of fruit

### SPRAYING INSTRUCTIONS

Use a hand sprayer that gives good coverage at sufficient pressure. When spraying underside of leaves a half pint per 100 sq ft is necessary. In case of large trees, it is necessary to spray before dusk and repeat sprayings at regular intervals. In case of lawns and other areas, a backpack sprayer may be used by following manufacturer's instructions.

### CAUTION

Do not breathe dust. Keep away from children. Do not use near water. Do not use near streams, ponds, lakes, or other bodies of water. Do not use near water when water is being used for drinking. Do not use near water when water is being used for drinking. Do not use near water when water is being used for drinking.