

VIGORO

50% CYTHION THE PREMIUM GRADE MALATHION SPRAY

ACTIVE INGREDIENTS	
Malathion*	50.0%
Heavy Aromatic Naphtha	42.0%
INERT INGREDIENTS	8.0%
	Total 100.0%

E P A Reg No 557-1887

E P A Est No 557-FL-1

*0,0-Dimethyl Dithiophosphate of Diethyl Mercapto-succinate

Store at temperatures above 32 F

Manufactured By Swift Chemical Company (An Estech Company) Chicago, Illinois 60664

NET CONTENTS 16 FL. OZ. (1 PT.)

CAUTION: Keep out of reach of children. See side panel for additional cautions.

KILLS:
APHIDS
CERTAIN SCALES
WHITEFLY
SPIDER MITES
MEALYBUGS
LEAFHOPPERS
THRIPS

DIRECTIONS FOR USE

Mix Vigoro 50% Cythion Spray with water in amounts indicated below and apply with a suitable sprayer. Cover plants and both sides of foliage thoroughly and repeat at 7 to 10 day intervals if necessary.

ORNAMENTALS: Azaleas, Camellias, Gardenias, Roses, Chrysanthemums, Cantons, Bougainvillea, Gladiolus, Hibiscus, Ixora, Philodendron, Evergreen, Red Spider Mites, Mealybugs, Woolly Aphids, White Fly, Thrips, Lace Bug, etc. ~~Apply 1 to 2 teaspoonful per gallon of water.~~
FRUITS: Citrus, Grapes, Peaches, etc. ~~Apply 1 to 2 teaspoonful per gallon of water.~~
VEGETABLES: Tomatoes, Beans, Peas, Cucumbers, Squash, Melons, Broccoli, Cabbage, Cauliflower, Kale, Mustard, Turnips, Eggplant, Peppers, Potatoes, Onions, Lettuce, Spinach, Celery, etc. ~~Apply 1 to 2 teaspoonful per gallon of water.~~

FRUITS: (Citrus, Grapes, Peaches) Scale crawlers and Aphids 1 1/2 to 2 teaspoonfuls per gallon of water. Red Spider Mite, Florida Red Scale, Soft Pink Scale, Purple Scale, Latania Scale, Plum Curculio, Mealybugs, White Fly, Thrips. Do not apply on Citrus, Avocado or Peaches within 7 days, on Grapes within 3 days, or on Mangoes within 2 days of harvest.

VEGETABLES: (Tomatoes, Beans, Peas, Cucumbers, Squash, Melons, Broccoli, Cabbage, Cauliflower, Kale, Mustard, Turnips, Eggplant, Peppers, Potatoes, Onions, Lettuce, Spinach, Celery) Aphids 1 1/2 to 2 teaspoonfuls per gallon of water. Red Spider Mites, Bean Leafhopper, Potato Leafhopper, Mexican Bean Beetle, Imported Cabbage Worm, Cabbage Looper, Mealybugs, White Fly, Thrips. 1 tablespoonful per gallon of water. Do not apply on Lettuce within 10 days, on Cabbage, Cauliflower, Kale, Mustard Greens, Spinach or Celery within 7 days, on Tomatoes, Beans, Peas, Broccoli, Turnips, Eggplant, Peppers, or Onions within 3 days, on Cucumbers, Squash, or Melons within 1 day of harvest.

CAUTION: Harmful if swallowed. Avoid breathing of vapor or spray mist. Avoid contact with skin. Wash thoroughly after using with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

This product is toxic to fish, shrimp and crab. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ANTIDOTE: Antidote if swallowed: Give a tablespoon of salt in a glass of warm water and repeat until vomit is clear. Have victim lie down and keep quiet. Call a physician immediately.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. Atropine is antidote.