


NET WT 30 LBS

# SWIFT MICRO-GRANO SULFUR (WETTABLE)



3021-1  
January 30, 1975  
UNDER THE FEDERAL  
567-1898

### DIRECTIONS FOR USE

CITRUS: Rust Mite—Use 6 pounds per 100 gallons of water in dilute sprays or equivalent ratio for concentrate applications.

### CAUTION

Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty drum or bag. If a drum, return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies. If a bag, destroy by burying or burning. Keep away from smoke and fumes.

Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality.

During period of high temperatures, sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures (over 90°F). Do not use sulphur with oil, or within 3 weeks after an oil application.

### ACTIVE INGREDIENT:

Sulphur, not less than .....95.0%

INERT INGREDIENTS: .....5.0%

Total 100.0%

E.P.A. Reg. No. 557-1898

E.P.A. Est. No. 557-FL-1

Particle Size—Average 5 Microns

**CAUTION:** Keep Out Of Reach Of Children.

See Back Panel For Additional Cautions.

Manufactured By  
Swift Chemical Company  
(An Estech Company)  
Chicago, Illinois 60604

NET WT 20 LBS

# SWIFT

## NUTRITIONAL SPRAY T-8 FOR CITRUS

2/12/75

557-1900

### DIRECTIONS FOR USE FOR FUNGUS DISEASES

Melanose and Scab - Use at rate of 4 pounds per 100 gallons of water, one to three weeks after petal fall. Where Melanose has been severe in the past or during wet springs, a second application should be made 4 weeks later. On varieties severely affected by Scab, make one application prior to spring flush of growth and another one to three weeks after petal fall.

Swift Nutritional Spray T-8 For Citrus contains basic copper sulfate, neutral zinc, manganese sulfate and neutral manganese. Used at the rate of 4 pounds per 100 gallons of water, the standard dosage of copper and manganese will be furnished. Do not apply this product separately as a nutritional spray - only when fungus disease is a problem.

### CONTAINS

### COPPER-ZINC-MANGANESE

#### ACTIVE INGREDIENT

Copper (in basic copper sulfate) as metallic ... 18.10%

#### INERT INGREDIENTS

81.90%  
Total 100.00%

#### SECONDARY PLANT FOODS

Zinc as ZnO ... 15.85%  
Manganese as MnO ... 17.60%

E P A Reg. No. 557-1900

E P A. Est. No. 557-FL-1

**CAUTION:** Keep Out Of Reach Of Children  
See Side Panel For Additional Cautions

### MIXING INSTRUCTIONS

Add to spray tank when 2/3 full of water and agitator is running. It is not necessary to add lime to this mixture for neutralization.

#### CAUTION

Harmful if swallowed. Wash hands, arms and face with soap and water after using. Avoid contamination of feed and foodstuffs.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

ANTIDOTE: Call a physician immediately. Give large quantities of whites of egg or milk.

Manufactured By  
Swift Chemical Company  
(An Estech Company)  
Chicago, Illinois 60604