

**FOR USE
ON
ORNAMENTAL
TURF
AND
LAWN GRASSES
ONLY**



**CUSTOM FORMULATED FERTILIZER
PLUS BENEFIN
FERTILIZER
12-6-8**

Minimum Guaranteed Analysis

ACTIVE INGREDIENT		
*N-butyl-N-ethyl-a, a-trifluoro-2, 6-dinitro-p-toluidine	0.75%	
INERT INGREDIENTS	99.25%	
E.P.A. Reg. No. 557-1919		TOTAL 100.00%
E.P.A. Est. No. 557-LL-1		

*Contains 15 pounds of technical Benefin per ton
**CAUTION: Keep Out Of Reach Of Children.
 See Below For Additional Cautions.**

Total Nitrogen	12.00%
Nitrate Nitrogen	2.10%
Ammoniacal Nitrogen	9.90%
Water Soluble Organic Nitrogen	None
Water Insoluble Nitrogen	None
Available Phosphoric Acid	6.00%
Water Soluble Potash	8.00%
Total Available Primary Plant Food	26.00%
Chlorine, not more than	1.00%

Primary Plant Food Sources: Sulphate of Ammonia, Triple Superphosphate, Ammonium Nitrate, Muriate of Potash, Sulphate of Potash, Magnesia, Sulphate of Potash.

SECONDARY PLANT FOODS	
Iron Sulphate as Fe ₂ O ₃	1.00%

Secondary Plant Food Sources: Iron Sulphate
 Manufactured by Swift Agricultural Chemicals Corporation, (An Estech Company), Chicago, Illinois 60604, Winter Haven and Pompano Beach, Florida, Fla. Reg. No. 732-735-5061.

DIRECTIONS FOR USE

For Southern Turf Grasses (Bermuda, Centipede, Fescue, Zoysia, St. Augustine, and Bahia) - To control Poa Annua, smooth crabgrass, hairy crabgrass, goose grass (silver crabgrass), water grass (barnyard grass), yellow and green foxtail. Apply broadcast at a rate 400 pounds per acre (9.0 pounds per 1000 square feet) prior to germination of weed grasses as the material will not control established weeds.

Do not apply to newly seeded areas. Do not apply in the spring to lawns planted the previous fall. Do not apply to newly sprigged areas until the turfs are well established. Do not mow within 48 hours after application.

CAUTION

Keep out of reach of children. Avoid contact with eyes. If contact with skin, or irritation, flush with water. Wash thoroughly after use. Do not breathe dust. Keep away from food and drink.

The product is toxic to fish. Keep out of lakes, streams, ponds, etc. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not treat or store in plastic whenever possible.

Apply this product to lawns as directed on the label.

Handwritten notes: 5/20/99, 10/1/99, 10/1/99



CUSTOM FORMULATED FERTILIZER PLUS BENEFIN 75

FOR USE ON ORNAMENTAL TURF AND LAWN GRASSES ONLY



CUSTOM FORMULATED FERTILIZER PLUS BENEFIN FERTILIZER 12-6-8

Minimum Guaranteed Analysis

ACTIVE INGREDIENT: *N-butyl-N-ethyl-a, a-trifluoro-2, 6-dinitro-p-toluidine ... 0.75%
INERT INGREDIENTS: 99.25%
E.P.A. Reg. No. 557-1919 TOTAL 100.00%
E.P.A. Est. No. 557-FL-1

*Contains 15 pounds of technical Benefin per ton.

CAUTION: Keep Out Of Reach Of Children. See Below For Additional Cautions.

Ammoniacal Nitrogen 2.10%
Water Soluble Organic Nitrogen 9.90%
Water Insoluble Nitrogen None
Available Phosphoric Acid 6.00%
Water Soluble Potash 8.00%
Total Available Primary Plant Food 26.00%
Chlorine, not more than 7.00%
Primary Plant Food Sources: Sulphate of Ammonia, Triple Superphosphate, Ammonium Nitrate, Muriate of Potash, Sulphate of Potash, Magnesia, Sulphate of Potash.

SECONDARY PLANT FOODS:
Iron Sulphate as Fe₂O₃ 1.00%

Secondary Plant Food Sources: Iron Sulphate
Manufactured by Swift Agricultural Chemicals Corporation, (An Estech Company), Chicago, Illinois 60604, Winter Haven and Pompano Beach, Florida, Fla. Reg. No. 732-735-5061.

DIRECTIONS FOR USE

For Southern Turf Grasses (Bermuda, Centipede, Fescue, Zoysia, St. Augustine, and Bahia) - To control Poa Annua, smooth crabgrass, hairy crabgrass, goose grass (silver crabgrass), water grass (barryard grass), yellow and green foxtail: Apply broadcast at a rate 400 pounds per acre (8.0 pounds per 1000 square feet) prior to germination of weed grasses as the material will not control established weeds.

Do not apply to newly seeded areas. Do not apply in the spring to lawns planted the previous fall. Do not apply to newly-sprigged areas until the turfs are well-established. Do not reseed within 42 days after application.

CAUTION

Keep out of reach of children. Avoid prolonged or repeated contact with skin, or inhalation of dust. Wash thoroughly after handling. Keep away from feed or foodstuffs.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse container. Destroy when empty.

Apply this product only as specified on this label.

