

SEP 27 1985

Mr. Bruce W. Michel
Estech, Inc.
30 North LaSalle Street
Chicago, IL 60602

Dear Mr. Michel:

Subject: Gro-Tone Liquid Sevin
EPA Registration No. 557-1944
Your Application Dated June 27, 1985

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

At the next printing of the label revise as follows:

The headings "Precautionary Statements," followed by "Hazards to Humans and Domestic Animals," "Environmental Hazards," "Physical and Chemical Hazards" must appear on the label followed by the appropriate wording. Refer to 40 CFR 162.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

BEST AVAILABLE COPY

RD/IRB:JOB:39419:Ellenberger:155776:RD-71:KENCO:9/26/85:del.10/4/85:dej

CONCURRENCES

SYMBOL	SURNAME	DATE					

9 2/2

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREE: Rose; Chrysanthemum, Arborvitae, Azalea, Juniper, Elm, Maple, Oak, Dogwood and Pine. To control: Japanese Beetles, Rose Aphids, Periodical Cicada, Mealy Bugs, Leafhoppers, Lace Bugs, Flea Beetles, Aphids, Bagworms, Rose Slugs, Scale Insects, Thrips (exposed). Apply 4 teaspoonfuls of Sevin insecticide per gallon of water, when insects or their damage appear. Completely cover all parts of plants, including underside of leaves and branches. Repeat application in 7 to 10 days as necessary.

LAWN INSECTS: Sod Webworms, Fall Armyworms, Fleas and Ants. Use 6 fluid ounces (1 cup) in 20 gallons water and cover 600 square feet of established lawn when insects first appear. Repeat in 2 or 3 weeks as needed. Keep children and animals off treated lawns until these areas are dry. For best results, mow lawn, remove clippings, and apply immediately after rain or watering. Do not water for at least 2 days after application.

VEGETABLES: Beans, Corn, Peppers, Tomatoes, and Potatoes. To control Potato Beetles on Potatoes and Tomatoes, Mexican Bean Beetles on Beans, Flea Beetles, Japanese Beetles on Beans and Corn, Armyworms, Tomato Hornworms, and Tomato Fruitworms on Tomatoes, Peppers and Potatoes. Mix 4 to 6 teaspoonful per gallon of water. Use 1 gallon of dilute spray to 430 square feet. Apply when insects or damage appear and repeat in 7 to 10 days as necessary. Potatoes - Foliage application only. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

E.P.A. Reg. No. 557-1944

Some injury may occur when used on tender foliage during prolonged wet periods or during periods of high humidity. Do not use on Boston Ivy.

PRECAUTIONARY INFORMATION: CAUTION! May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Keep container closed. Use with adequate ventilation. Avoid storage near feed and food products. Wash thoroughly after handling.

FIRST AID: If swallowed, induce vomiting by sticking finger down throat. Repeat until vomit is clear. Call physician. Never give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiate or cholinesterase inhibiting drugs.

IMPORTANT: Shake well before using. Agitate during mixing and spraying. Keep container closed to prevent evaporation. Do not store below 32° F.

ENVIRONMENTAL PRECAUTIONS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONTAINER DISPOSAL: Do not reuse container. Destroy when empty.

CONDITIONS OF SALE: All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.

EPA Est No 2217-KS-1

BEST AVAILABLE COPY

Under
Fur
ns are
register

GR

Liquic

CO

Ants, Aphids, Bagworms,
Japanese Beetles, Lace
Mexican Bean Beetles, Pe
Aphids, Rose Slugs, Sc
(exposed), Tomato Fruitw

APPLY TO

ACTIVE INGREDIENT
Carbaryl (1-naphthyl N-m
INERT INGREDIENTS
†Seven insecticide, Sevin,
Corporation.

KEEP OUT OF

See back panel for addi
specific directions for use

Net

Distributed by ESTECH, INC.
30 North LaSalle Street
Chicago, Illinois 60602