



CERTIFICATION OF COMPLIANCE

I, Judith K. Lucash, being an authorized representative of Vigoro Industries, Inc., certify that all containers of

Par Ex Slow Release Fertilizer

EPA Reg. No. 4557-1952

produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

Judith K. Lucash
Manager of Regulatory Affairs

December 21, 1992

2075

FRONT PANEL

TREATS 15,000 SQ.FT.
26-3-5

SLOW RELEASE FERTILIZER

WEED & FEED
WITH IBDU/SCU®

PAR EX®

PROFESSIONAL PRODUCTS

Provides uniform continuous feeding for your lawn as it kills Chickweed, Dandelion, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds.

ACTIVE INGREDIENTS:

*2,4-dichlorophenoxyacetic acid	0.66%
2-(2-methyl-4-chlorophenoxy) propionic acid	0.30%
Dicamba (3,6-dichloro-o-anisic acid).	0.06%
INERT INGREDIENTS:	98.98%

TOTAL 100.00%

*Isomer Specific by AOAC Method 6.D01.5

For sale to, use and storage by professional turf maintenance personnel only.

KEEP OUT OF REACH OF CHILDREN


CAUTION

In case of contact: for eyes, flush with water for 15 minutes; for skin, wash with plenty of soap and water. Get medical attention if irritation persists. See back panel for additional precautionary statements.

EPA Reg. No. 557-1952
U.S. Patent No. 3,284,186

EPA Est. No. 557-CT-1

NET WT. 50 LBS.



NET WT. 50 LBS.

26-3-5

slow release fertilizer
WEED & FEED
WITH IBDU/SCU®

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	26.0%
1.2% Ammoniacal Nitrogen	
1.5% Water Insoluble Nitrogen	
5.0% Coated Slow Release Urea Nitrogen	
18.3% Urea Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	3.0%
SOLUBLE POTASH (K ₂ O)	5.0%

Derived from isobutylidene diurea, sulfur-coated urea, urea, ammoniated phosphate and muriate of potash. Potential Acidity 600 lbs. Calcium Carbonate Equivalent per ton.

PAR EX®

PROFESSIONAL PRODUCTS

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes and clothing. Avoid breathing dust. Wash thoroughly after handling. Harmful if swallowed. Keep out of reach of children.

Protective Clothing: When using this product, wear long-sleeved shirt, long pants, socks, shoes and gloves. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARD

This product is toxic to aquatic invertebrates. Drift or run-off may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

REAR PANEL Cont.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment areas during application. Do not enter treatment areas until spray has dried or dust has settled.

THREE WAY ACTION - in a single application PAR EX® Slow Release Fertilizer Weed and Feed provides uniform continuous feeding for your lawn while it kills many broadleaf weeds by contact and systemic action. For best results, apply when weeds are actively growing and when soil moisture is adequate. If adverse low moisture conditions exist, weeds are in a poor state of growth, or area is heavily overrun with weeds, retreatment may be necessary. make a second application in approximately 30 days. This product may be used on bluegrass, fescue, rye, Bermuda and zoysia lawns.

1. Apply contents of this bag to 15,000 sq. ft.
2. Mow lawn to normal height 1 or 2 days before application.
3. Water lawn thoroughly 1 day before application to ensure adequate soil moisture.
4. Do not water lawn again until 1 day after application.

The maximum application rate for this product is 3.3 lbs. per 1,000 sq. ft. The maximum number of broadcast applications per treatment site is 2 per year.

Approximately 4 weeks after weed kill is evident, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

When applied as directed, the following weeds in established turfgrasses are controlled.

Bedstraw	Knotweed	Sheep Sorrel
Black Medic	Lambsquarter	Shepherd's Purse
Buckhorn	Lespedeza	Spoodwell
Burdock	Morning Glory	Spurge
Chicory	Oxalis	Thistle
Chickweed	Peppergrass	Wild Carrot
Clover	Pigweed	Wild Garlic
Dandelion	Plantains	Wild Lettuce
Dock	Poison Ivy	Wild Onion
Ground Ivy	Poison Oak	Yarrow
Heal-all	Purslane	and many other
Henbit	Ragweed	broadleaf weeds

5045

REAR PANEL Cont.

STORAGE AND DISPOSAL

STORAGE: Store in a dry place. Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities by burning. If burned, stay out of smoke.

PRECAUTIONS - Do not apply to bentgrass, carpetgrass, St. Augustine grass or Dichondra lawns, as injury may result. Do not use where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. Apply when air is calm. Do not apply to areas underlaid by roots of desirable trees and shrubs.

SUGGESTED SPREADER SETTINGS

SPREADER	WIDTH	SETTING
Cyclone	9 ft.	4
Central "Roto-Spred"	10 ft.	3
Scott PF-1		5 1/2
Scott R7 Professional	10 ft.	G
Lely - HR 425 rpm 4 mph	30 3/4 ft.	5 1/2
Lely - WRT @ 4.5 mph		4 3/4

Note: Spreader settings are approximate. Check by trial over known area.

CAUTION: Close spreader openings when loading, turning or when stopped.

Manufactured by Vigoro Industries, Inc., Fairview Heights, IL 62208. PAR EX® and IBDU/SCU® are registered trademarks of Vigoro Industries, Inc.

