

In Accordance with the...  
Registration No. 111-111-1111



**PLUS WEED CONTROL**

**pursley LAWN FOOD**

**PURSLEY  
LAWN FOOD  
PLUS  
WEED CONTROL**

- ✓ Covers 3,000 sq. ft.
- ✓ Formulated especially for Seville, Floratam, Fioratine, Bitterblue, Zoysia and Centipede grasses.
- ✓ Contains slow release nitrogen, iron, manganese and sulphur
- ✓ Contains Atrazine for weed control
- ✓ Long-lasting

KEEP OUT OF REACH OF CHILDREN  
CAUTION

**PURSLEY LAWN FOOD PLUS WEED CONTROL**

# PURSLEY LAWN FOOD PLUS WEED CONTROL

Contents Treats 3 000 Sq Ft Apply 10 lbs per 1000 Sq Ft (20 x 50 Area)

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

### ENVIRONMENTAL HAZARD

Atrazine can leach through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable (i.e. well-drained). Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**Established Lawns:** For best results apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in spring and fall. Apply when lawn is wet with dew or immediately following rain or watering. Wait 24 hours before watering in.

**New Lawns:** Apply immediately after planting to new lawns that have been plugged or sprigged to prevent growth of weeds that come from seed. Water in well.

Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass or carpetgrass or where clover is present and desired.

Avoid application to the root zones of trees and shrubs.

Avoid applications on garden plants, flowers and shrubs.

Delay spring planting for one month after application. Do not apply in areas underlaid by roots of desirable trees and shrubs. Clean spreader after use. Apply only in Winter, Spring and Fall. This product is not recommended for use during the months of July and August.

Use herbicides properly. Do not over apply. Read directions for use carefully. Water in as directed. Apply spring application between January 1 and March 31 in Florida and other southern states to June 15th.

### CONTROLS THESE WEEDS

Germinating Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings, Dallisgrass, Vaseygrass, Bahiagrass, Sedges (such as Cyperus Compressus) and Sours.

Established and Germinating: Betony, Black medic (yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pursley, Pennywort (Dollarweed) and Creeping Charlie (Matchweed).

### SPREADER SETTING

Spreader	Setting	Spreader	Setting
Gro-Tone Rotary	6	Sears Mark II	8
Vigoro Rotary	6	Wards	5
Vigoro Drop	7	Lawn Beauty	5
Cyclone	6	Scotts PF-1	7

NOTE: Spreader rate varies with walking speed; check spreader over known area.

**STORAGE:** Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

**PRODUCT DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

### GUARANTEED ANALYSIS

735-4576

Total Nitrogen (N)		12.00%
Water Soluble Organic Nitrogen	7.50%	
Water Insoluble Nitrogen	4.50%	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )		3.00%
Soluble Potash (K <sub>2</sub> O)		3.00%
Chlorine, not more than		2.00%
Derived from Tankage, Isobutylidene Diurea, Urea, Triple Superphosphate, Sulphate of Potash, Sulphate of Potash Magnesia		
Secondary Nutrients		
Total Magnesium (Mg)		0.50%
Water Soluble Magnesium (Mg)		0.50%
Sulphur, combined (S)		1.40%
Boron (B)		0.02%
Copper (Cu)		0.07%
Iron (Fe)		2.00%
Chelated Iron (Fe)		0.02%
Manganese (Mn)		1.00%
Zinc (Zn)		0.08%
Derived from Sulphate of Potash Magnesia, Sulphate of Potash, Sodium Borate, Copper Oxide, Copper Sulphate, Iron Oxide, Iron Sulphate, Chelated Iron EDTA-HEEDTA, Manganous Oxide, Manganese Sulphate, Zinc Oxide and Zinc Sulphate		

### CAUTION

Keep out of reach of children. Most fertilizers may stain paths and sidewalks. Remove spillage promptly.

Manufactured by VIGORO INDUSTRIES INC. FAIRVIEW HEIGHTS, IL 62208

PURSLEY LAWN FOOD PLUS WEED CONTROL