

P U R S L E Y
L A W N F O O D
P L U S
W E E D C O N T R O L

P U R S L E Y
L A W N F O O D
P L U S W E E D
C O N T R O L

- Covers 3,000 sq. ft.
- Formulated especially for Seville, Floratam, Floratine, Bitterblue, Zoysia and Centipede grasses.
- Contains slow release nitrogen, iron, manganese and sulphur.
- Contains Atrazine for Weed control.
- Long-lasting.

ACCEPTED
APR 30 1990
United States Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 557-1953

P U R S L E Y L A W N F O O D P L U S W E E D C O N T R O L

KEEP OUT OF REACH OF CHILDREN
C A U T I O N
In case of contact, for skin, wash with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention if irritation persists. See back panel for additional precautionary statements.

ACTIVE INGREDIENT

*Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine).....	0.40%
*Related compounds	0.02%
INERT INGREDIENTS	99.58%
TOTAL	100.00%

*Contains 8.4 lbs. of atrazine and related compounds per ton.

E.P.A. Reg. No. 557-1953
E.P.A. Est. No. 46930-FL-1

Net Wt. 30 lbs.
(13.63 Kg)

BEST AVAILABLE COPY

NET WT. 30 LBS. (13.63 Kg)

PURSLEY
LAWN FOOD PLUS
WEED CONTROL

Contents Treats 3,000 Sq. Ft. Apply 10 lbs. per 1,000 Sq. Ft. (20'x50' Area)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

C A U T I O N

Avoid contact with skin, eyes or clothing
Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

Atrazine can leach through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Established Lawns: For best results apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in spring and fall. Apply when lawn is wet with dew or immediately following rain or watering. Wait 24 hours before watering in.

New Lawns: Apply immediately after planting to new lawns that have been plugged or sprigged to prevent growth of weeds that come from seed. Water in well.

Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass, or carpetgrass or where clover is present and desired.

Avoid applications to the root zones of trees and shrubs.

Avoid applications on garden plants, flowers and shrubs.

Delay spring planting for one month after application. Do not apply in areas underlaid by roots of desirable trees and shrubs. Clean spreader after use. Apply only in Winter, Spring or Fall. This product is not recommended for use during the months of July and August.

BEST AVAILABLE COPY

REAR PANEL CONT.

Use herbicides properly. Do not over-apply. Read directions for use carefully. Water in as directed. Apply spring application between January 1 and March 31 (Florida and other southern states to June 15).

Approximately 4 weeks after initial treatment with product, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

CONTROLS THESE WEEDS

Germinating: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings (Dallisgrass, Vaseygrass, Bahiagrass), Sedges (such as Cyperus Compressus) and Spurge.

Established and Germinating: Betony, Blackmedic (yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort (Dollarweed) and Creeping Charlie (Matchweed).

SPREADER SETTINGS

SPREADER	SETTING	SPREADER	SETTING
Gro-Tone Rotary	6 1/4	Sears Mark II	8 3/4
Vigoro Rotary	6 1/4	Wards	5 1/4
Vigoro Drop	7 1/4	Lawns Beauty.	5
Cyclone	6 1/4	Scotts PF-1	7

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not re-use bag. Discard bag in trash.

ulu

GUARANTEED ANALYSIS

735-4576

TOTAL NITROGEN (N)	12.00%
7.50% Urea Nitrogen	
4.50% Water Insoluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	3.00%
SOLUBLE POTASH (K ₂ O)	3.00%
Chlorine, not more than	2.00%
Derived from Tankage, Isobutylidene Diurea, Urea, Triple Super-phosphate, Sulphate of Potash, Sulphate of Potash Magnesia.	

Statement of Secondary Nutrients

Total Magnesium (Mg)	0.50%
0.50% Water Soluble Magnesium (Mg)	
Sulphur, combined (S)	1.40%
Boron (B)	0.02%
Copper (Cu)	0.07%
Iron (Fe)	2.00%
Iron, Chelated (Fe)	0.02%
Total Manganese (Mn)	1.00%
Zinc (Zn)	0.06%
Derived from sulphate of potash magnesia, sulphate of potash, manganous oxide, copper oxide, copper, sulphate, zinc oxide, zinc sulphate, sodium borate, iron oxide, iron sulphate and iron sulphate EDTA-HEEDTA.	

CAUTION

Keep out of reach of children. Most fertilizers may stain patios and sidewalks. Remove spillage promptly.

Manufactured by Vigoro Industries, Inc., Fairview Heights, IL 62208.