

NOT REVIEWED
In Accordance with PR Notice 82-11
Based on Draft Labeling Dated 3/8/85

PARK RIDGE

PARK RIDGE 20

WEED & FEED

25-3-3 WEED & FEED

Essex Inc
an Emerson Company

25-3-3

Weed control with lawn fertilizer
Apply when weeds are growing
Controls dandelions and broadleaf weeds
Lawn fertilizer provides thick green lawns

WENT OUT OF RANGE OF CONTROL

CAUTION

In case of contact, wash skin with plenty of soap and water for 15 min. Flush with water for 15 minutes and get medical attention. See back panel for additional precautionary measures.

ACTIVE SUBSTANCES	
*Benzylamine Salt of 2-(2-methyl-4- chlorophenyl) propionic acid	0.540%
*Benzylamine Salt of 2,4-dichlorophenoxyacetic acid	0.140%
*Benzylamine Salt of dimethyl (1,3-dichloro-5-oxo) acid	0.007%
WORTH September 1972	92.200%

NEW PRODUCT CONTINUED

- **20PP Acid** - 0.47% - 0.74 lbs. per acre on 0.50 lbs. per ton
- **24-D Acid** - 0.17% - 0.21 lbs. per acre on 2.40 lbs. per ton
- **Dicamba Acid** - 0.21% - 0.40 lbs. per acre on 1.90 lbs. per ton
- **Inorganic Sulfuric L, ADRC Mixture 6 (D1) 5**


U.S. Patent No. 3,201,000
©1974 Dow Co. 047-1000 047-1000 lbs. 047-1

Part Master™ is a trademark of Entech, Inc.

NET WT 20 LBS. (9.1Kg)

BEST AVAILABLE COPY

25-3-3
WEED & FEED



PARK
RIDGE

PARK RIDGE™ WEED & FEED 25-3-3

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	25.0%
1.0% Ammoniacal Nitrogen	
5.0% Coated Slow Release Urea Nitrogen	
19.0% Urea Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	3.0%
SOLUBLE POTASH (K ₂ O)	3.0%
Chlorine, not more than	3.0%

Derived from ammoniated phosphates, ammonium sulfate, sulfur-coated urea and muriate of potash. Potential Acidity 500 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Estech, Inc., Chicago, Illinois 60602

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing dust.
Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not apply when weather conditions favor drift from target area.
Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Mow lawn to normal height 1 or 2 days before application.
2. Water lawn thoroughly 1 day before application to ensure adequate soil moisture.
3. Apply contents of this bag to 5,000 sq. ft. using the suggested spreader setting. **DO NOT OVERAPPLY**. For best results, apply when weeds are actively growing.
4. Do not water lawn again until 1 day after application.

This product may be used on St. Augustine, Centipede, Bahia, Bermuda, Fescue, Zoysia and Bentgrass lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or area is heavily overrun with weeds, retreatment may be necessary. Make a second application in approximately 30 days.

This product combines fast acting and long-lasting feeding - plus controls the following weeds:

Florida Pusley	Knotweed	Ragweed
Black Medic	Lambsquarter	Sida
Chickweed	Mustard	Spurge
Clover	Orals	Wild Onion
Cocklebur	Peppergass	Wild Lettuce
Dandelion	Pigweed	and many other
Creeping Charlie	Plantains	Broadleaf
Dock	Poison Ivy	Weeds
Henbit	Purslane	

PRECAUTIONS: - Do not apply to newly seeded lawns. Do not apply to dichondra, lippie or other ground cover. Avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly because temporary foliage discoloration may occur. Do not use where desirable clovers are present or allow the product to "drift" on flowers, vegetables or ornamental shrubbery. Apply when air is calm. Do not apply to areas underlaid by roots of desirable trees and shrubs. During application, close spreader openings when turning.

SUGGESTED SPREADER SETTINGS

SPREADER	SETTING	SPREADER	SETTING
Wards 1091, 1093, 1095	6%	Precision 20C Drop	4
Wards 1080, 1081	5	SB 40 Broadcast	12 1/2
Jackson & Sears	5	24F Drop	1
Scott	5 1/2	Central 40 Drop	5%
Kmart	5 1/2	60 Broadcast	6%
		Cyclone	4%

NOTE: Spreader rate varies with walking speed. To insure proper coverage, check spreader over known area.

STORAGE: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, feed or food. Return spillage to original container prior to disposal.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.
CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

NET WT. 20 LBS. (9.1Kg) TREATS 5,000 SQ. FT.