PARK RIDGE 18

PARK RIDGE

# WEED & FEED

25-3-3 WEED & FEED

TREATS 5,000 SQ. FT.

25-33

Weed control with lawn fertilizer Apply when weeds are growing Controls dandelions and broadleaf weeds Lawn fertilizer provides thick green lawns

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF THE ACT OF CHILDREN
CAUTION

\*\*TOTAL PROPERTY OF CHILDREN
CAUTION
CAUTION

\*\*TOTAL PROPERTY OF CHILDREN
CAUTION

NET WT. 18 LBS. (8.1 Kg)



WEED & FEED

# **PRECAUTIONARY STATEMENTS** HAZARD TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

#### ENVIRONMENTAL HAZARD

 $(x_1, x_2, \dots, x_n)$  , where  $(x_1, \dots, x_n)$  , we see that  $(x_1, \dots, x_n)$  , where  $(x_1, \dots, x_n)$  , where  $(x_1, \dots, x_n)$ 

#### DIRECTIONS FOR USE

#### SUGGESTED SPREADER SETTINGS

	• · • • • · ·		
SPREACHER	SETTING	SPREADER	SETTING
		the second second	:
••	• •	4.00.468.4	
5 P 2	4 %	21	
4 - 1 No. 4 4	3 %	A CAR A STATE OF THE STATE OF T	
		egic in the desired	: •
•			:

PROTECTION AND ACTION AND ACTION AND ACTION AND ACTION AND ACTION AND ACTION ASSESSMENT AND ACTION ASSESSMENT AND ACTION ASSESSMENT ASSESSMENT

STORMAGE of entringing out performs outputs to a control of the co

NET WT. 18 LBS. (8.1 Kg) TREATS 5,000 SQ. FT.

FRONT PANEL

557 1959

## PARK RIDGE® 18

WEED & FEED 25-3-3

Treats 5,000 sq ft

Weed Control with lawn fertilizer - Apply when weeds are growing.

Controls dandelions and broadleaf weeds Lawn fertilizer provides thick green lawns.

#### ACTIVE INGREDIENTS

**1	Dimethylamine	Salt o	f 2-(2-methyl-4-chlorophenoxy)propionic acid	0.567%
+^; ***!	Dimethylamine	Salt o	f 2,4-dichlorophenoxyacetic acid	0.163%
~~~;	Jimetnylamine	zarr o	f dicamba(3,6-dichloro-o-anisic acid)	0.069%
INERT	INGREDIENTS		· · · · · · · · · · · · · · · · · · ·	99.201%
			TOTAL	100.00%

#### THIS PRODUCT CONTAINS:

\*\*MCPP Acid - 0.473% - 0.74 lbs. per acre or 9.46 lbs. per ton +\*2,4-D Acid - 0.136% - 0.21 lbs. per acre or 2.72 lbs. per ton \*\*\*Dicamba Acid- 0.058% - 0.09 lbs. per acre or 1.16 lbs. per ton


+Isomer Specific by AOAC Method 6.DO1-5

U.S.Patent No. 3,284,186 EPA Reg. No. 557-1959 EPA Est. No. 557-IL-1

# KEEP OUT OF REACH OF CHILDREN

## CAUTION

In case of contact, wash skin with plenty of soap and water: for eyes, flush with water for 15 minutes and get medical attention. See back panel for additional precautionary statements.



NET WT. 18 LBS. (8.1 Kg)

#### REAR PANEL

#### PARK RIDGE® 18

#### WEED & FEED

Treats 5,000 sq. ft. NET WT 18 lbs (8.1 Kg)

#### WITH LAWN FERTILIZER

25-3-3

## GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	25.0%
1.0% Ammoniacal Nitrogen 5.0% Coated Slow Release Urea Nitrogen 19.0% Urea Nitrogen	
AVAILABLE PHOSPHORIC ACID (P205)	3.0%
SOLUBLE POTASH (K <sub>2</sub> O)	
Chlorine, not more than	3.0%

Derived from ammoniated phosphates, ammonium sulfate, sulfur-coated urea and muriate of potash. Potential Acidity 500 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Estech, Inc., Chicago, Illinois 60602.

#### PRECAUTIONARY STATEMENTS

## HAZARD TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed. Avoid breathing dost. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

#### ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- 1. Mow lawn to nor al height 1 or 2 days before application.
- 2. Water lawn thoroughly 1 day before application to ensure adequate soil moisture.
- 3. Apply contents of this bag to 5,000 sq. ft. using the suggested spreader setting. DO NOT OVERAPPLY. For best results, apply when weeds are actively growing.
- 4. Do not water lawn again until I day after application.

This product may be used on St. Augustine, Cantipede, Bahia, Bermuda, Fescue, Zoysia and Bentgrass lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or area is heavily overrun with weeds, retreatment may be necessary. Make a scoond application in approximately 30 days.

This product combines fast-acting and long-lasting feeding - plus controls the following weeds: -

Florida Pusley
Black Medic
Chickweed
Clover
Cocklebur
Dandelion
Creeping Charlie
Dock

Henbit
Knotweed
Lambsquarter
Mustard
Oxalis
Peppergrass
Pigweed
Plantains

Poison Ivy
Purslane
Ragweed
Sida
Spurge
Wild Onion
Wild Lettuce
and many other
Broadleaf Weeds

PRECAUTIONS: - Do not apply to newly seeded lawns. Do not apply to dichondra, lippia or other ground cover. Avoid treating St. Augustine grass in summer months (June through Septemper) when it is growing rapidly because temporary foliage discoloration may occur. Do not use where desirable clovers are present or allow the product to 'wind drift' on flowers, vegetables or ornamental shrubbery. Apply when air is calm. Do not apply to areas underlaid by roots of desirable trees and shrubs. During application, close spreader openings when turning.

# SUGGESTED SPREADER SETTINGS

Spreader	Setting	Spreader		Setting
Wards 1091, 1093, 1095	4-3/4 4-3/4 5	SB 24F Central 40 60	Drop	12 1 5-1/4

NOTE: Spreader rate varies with walking speed. To insure proper coverage, check spreader over known area.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container isposal: Do not re-use bag. Discard bag in trash.

Park Ridge $^{\oplus}$  is a registered trademark of Estech, Inc.