

Reg # 557-1959

PM-23

192

NOT REVIEWED

In Accordance with PR Notice 82-2.

Based on Draft Labeling Dated 3/8/85

PARK
RIDGEPARK
RIDGE
18

WEED & FEED

25-3-3
WEED & FEED

TREATS 5,000 SQ FT

25-3-3

Weed control with lawn fertilizer
Apply when weeds are growingControls dandelions and broadleaf weeds
Lawn fertilizer provides thick green lawns

KEEP OUT OF REACH OF CHILDREN

CAUTION

In case of contact, wash skin with plenty of water and
water for 15 min. Flush with water for 30 minutes and get
medical attention. See label card for additional product
safety statements.

ACTIVE INGREDIENTS

2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt)	1.7%
2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt)	0.65%
2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt)	0.00%
2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt)	99.29%
INERT INGREDIENTS	
Total	100.00%

THIS PRODUCT CONTAINS

2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt)	0.46 lbs. per acre or 9.46 lbs. per ton
2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt)	0.21 lbs. per acre or 2.72 lbs. per ton
2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt)	0.00 lbs. per acre or 1.14 lbs. per ton
2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt)	0.00 lbs. per acre or 0.00 lbs. per ton

2,4-Dichlorophenoxyacetic acid (2,4-D) (as salt) is a Restricted Use Pesticide (RUP) under FIFRA. See label for use restrictions.

Eaton, Inc.
an Ecolab Company

Park Ridge™ is a trademark of Eaton, Inc.

NET WT. 18 LBS. (8.1 Kg)

**25-3-3
WEED & FEED**



**PARK
RIDGE**

PARK RIDGE™ WEED & FEED 25-3-3

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	25.0%
1.0% Ammoniacal Nitrogen	
5.0% Coated Slow Release Urea Nitrogen	
19.0% Urea Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	3.0%
SOLUBLE POTASH (K ₂ O)	3.0%
Chlorine, not more than	3.0%
Derived from ammoniated phosphates, ammonium sulfate, sulfur-coated urea and muriate of potash. Potential Acidity 500 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Estech, Inc., Chicago, Illinois 60602	

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing dust.
Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not apply when weather conditions favor drift from target area.
Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Mow lawn to normal height 1 or 2 days before application.
2. Water lawn thoroughly 1 day before application to ensure adequate soil moisture.
3. Apply contents of this bag to 5,000 sq. ft. using the suggested spreader setting. **DO NOT OVERAPPLY.** For best results, apply when weeds are actively growing.
4. Do not water lawn again until 1 day after application.

This product may be used on St. Augustine, Centipede, Bahia, Bermuda, Fescue, Zoysia and Bentgrass lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or area is heavily overrun with weeds, retreatment may be necessary. Make a second application in approximately 30 days.

This product combines fast-acting and long-lasting feeding - plus controls the following weeds:

Florida Pusley	Knotweed	Ragweed
Black Medic	Lambsquarter	Sida
Chickweed	Mustard	Spurge
Clover	Osage	Wild Onion
Cocklebur	Peppergass	Wild Lettuce
Dandelion	Pigweed	and many other
Creeping Charlie	Plantain	Broadleaf
Dock	Poison Ivy	Weeds
Henbit	Purslane	

PRECAUTIONS: - Do not apply to newly seeded lawns. Do not apply to dichondra, lippia or other ground cover. Avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly because temporary foliage discoloration may occur. Do not use where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables or ornamental shrubbery. Apply when air is calm. Do not apply to areas underlaid by roots of desirable trees and shrubs. During application, close spreader openings when turning.

SUGGESTED SPREADER SETTINGS

SPREADER	SETTING	SPREADER	SETTING
Wards 1091, 1093		Precision 20C Drop	4
1095	6%	SB 40 Broadcast	12
Wards 1080, 1031	4%	24F Drop	1
Jackson & Sears	4%	Central 40 Drop	5%
Scott	5	60 Broadcast	4%
Kmart	5	Cyclone	4

NOTE: Spreader rate varies with walking speed. To insure proper coverage, check spreader over known area.

STORAGE: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizers, seeds, food or feed. Return spillage to original container prior to disposal.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

NET WT. 18 LBS. (8.1 Kg) TREATS 5,000 SQ. FT.