

FRONT PANEL

Treats 5,000 sq. ft.


VIGORO CRABGRASS PREVENTER

PLUS LAWN FOOD

NORTHERN FORMULA

Ends the Crabgrass Menace

Feeds Your Lawn



KEEP OUT OF REACH OF CHILDREN

CAUTION

In case of contact; for eyes, flush with water for 15 minutes; for skin, wash with plenty of soap and water. Get medical attention if irritation persists. See back panel for additional precautionary statements.

ACTIVE INGREDIENT

N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine	0.87%
---	-------

INERT INGREDIENTS

99.13%

Total 100.00%

Net Wt. 20 lbs.

E.P.A. Reg. No. 557-1960

E.P.A. Est. No. 557-IL-1

REAR PANEL

NET WT. 20 LBS.

Treats 5,000 sq. ft.

VIGORO

CRABGRASS PREVENTER PLUS LAWN FOOD

NORTHERN FORMULA

15-3-3

TOTAL NITROGEN (N) 15.0%
11.7% Ammoniacal Nitrogen
0.3% Urea Nitrogen
3.0% Coated Slow Release Urea Nitrogen

AVAILABLE PHOSPHORIC ACID (P₂O₅) 3.0%
SOLUBLE POTASH (K₂O) 3.0%

Derived from ammonium sulfate, ammoniated phosphates, sulfur-coated urea and muriate of potash. Potential Acidity 800 lbs. Calcium Carbonate Equivalent per ton. Manufactured by Vigoro Industries, Inc., Fairview Heights, Illinois 62208. Vigoro is a registered trademark of Vigoro Industries, Inc.

KEEP OUT OF REACH OF CHILDREN
PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Remove contaminated clothing and wash before re-use.

Statement of Practical Treatment

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention.

REAR PANEL (contd.)

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not contaminate water by disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.
Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.
Container Disposal: Do not re-use bag. Discard bag in trash.

Apply Vigoro Crabgrass Preventer Plus Lawn Food early in the spring. It kills crabgrass and weed seedlings just as they are sprouting from the seed, so get it on early. This should be done one to two weeks prior to the germination of annual grass weeds. It will not control established weeds.

1. Rake area free of leaves and other debris prior to application.
2. Apply contents of this bag to 5,000 sq. ft.
3. It is essential to apply with no skips or misses. Calibrate and check equipment prior to use. Set spreader according to the chart below and apply evenly to the area. Check frequently to ensure equipment is working properly and distributing product uniformly.
4. Water in after application for optimum weed control.
5. Approximately 4 weeks after initial treatment with product, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft.

When applied as directed, the following weeds in established turf are controlled: crabgrass, foxtails and barnyard grass with an application in spring; goosegrasses with an early summer application. Vigoro Crabgrass Preventer Plus Lawn Food is tolerant to the following turf grasses when applied as directed:

Kentucky Bluegrass Perennial Ryegrass Fescue Zoysia

REAR PANEL (contd.)

PRECAUTIONS: This product may thin established annual bluegrass and should not be applied in the spring to turfgrass planted the previous fall. Do not apply to bentgrass putting greens or dichondra turf. Avoid any cultural practices that disturb the soil such as raking, aerification or verticutting after application of this product. Reseeding of treated areas should be delayed for at least 6 weeks after applying this product.

SUGGESTED SPREADER SETTING

SPREADER	SETTING	SPREADER	SETTING
Cyclone	4 1/8	Wards	5
Sears Mark II	5	Precision 20C	4
Central 40 Drop	5 3/4	Scotts PF-1	5 1/2
Central 60 Broadcast	4 3/8	Kmart	5 1/2

NOTE: Spreader setting varies with terrain, equipment and walking speed. To insure proper coverage, check spreader over known area.

CAUTION: Close spreader openings when loading, turning or when stopped.