



Treats 3000 Sq. Ft.

NET WT. 22 Lbs. 8 Oz.

HALL'S
WEED & FEED
FOR ST. AUGUSTINE

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of water.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For best results apply before weeds come up (preemergent). Established weeds are more difficult to control and may require an application in Spring and Fall. On newly sprigged St. Augustine, Centipede or Zoysia Lawns, apply immediately after planting to prevent weeds from retarding growth. Use herbicides properly.

Do not overapply. Read directions for use carefully. Water in well. Apply between January 1 and March 31 (Florida and other southern states to June 15th).

Apply $7\frac{1}{2}$ lbs. per 1000 sq. ft. (20 ft. x 50 ft. area) Contents of bag treats 3000 sq. ft.

CONTROLS THESE WEEDS

Germinating: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings (Dallisgrass, Vaseygrass, Bahiagrass). Sedges (such as Cyperus Compressus), Spurge and Sandburs.

Established and Germinating: Betony, Blackmedic (yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort (Dollarweed) and Creeping Charlie (Matchweed).

Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass, or carpetgrass or where clover is present and desired.

Avoid application to the root zones of trees and shrubs.

Avoid applications on garden plants, flowers and shrubs.

Delay spring planting for one month after application. Do not apply in areas underlaid by roots of desirable trees and shrubs. Clean spreader after use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during the months of July and August.

SPREADER SETTING

<u>Spreader</u>	<u>Setting</u>
Cyclone	4½
Scotts PF-1	4

Note: Spreader rate varies with walking speed, check spreader over known area.

STORAGE AND DISPOSAL

STORAGE-Keep away from feed or food products. Store in a dry place. Do not store near growing plants or fertilizers, seeds, insecticides or fungicides

DISPOSAL-Do not reuse empty container. Wrap container and put in trash collection.

GUARANTEED ANALYSIS

Reg. No. 731-8125

Total Nitrogen (N)-----	21.00%
Ammoniacal Nitrogen -----	1.56%
Water Soluble Organic Nitrogen -----	19.44%
Available Phosphoric Acid (P ₂ O ₅)-----	4.00%
Soluble Potash (K ₂ O)-----	4.00%
Chlorine, not more than -----	2.00%
Derived from Urea, DiAmmonium Phosphate, Triple Superphosphate, Sulphate of Potash.	

Statement of Secondary Plant Nutrients:

 Iron as Fe ----- 1.50%

Derived from Fritted Iron.

CAUTION

Keep out of reach of children. Most fertilizers may stain patios and sidewalks. Remove spillage promptly.

Manufactured by ESTECH, INC., Chicago, Illinois 60602