

Reg No: 557-1965

PM-25


194

HALL'S

NURSERIES & HARDWARES

WEED & FEED

FOR ST. AUGUSTINE



Use on St. Augustine, Centipede and Zoysia Grasses only.

1. Controls germinating weeds, including sandburs and certain established weeds as listed in the directions for use.
2. A complete plant food fortified with iron.
3. Covers 3,000 sq. ft.

KEEP OUT OF REACH OF CHILDREN

CAUTION

In case of contact, for skin, wash with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention if irritation persists. See back panel for additional precautionary statements.

ACTIVE INGREDIENT

*Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine .....	0.60
Related compounds .....	0.02
INERT INGREDIENTS .....	99.37
Total .....	100.00

\*Contains 12.60 lbs. of atrazine and related compounds per ton.

EPA Reg. No. 557-1965

EPA Est. No. 557-FL-

THREE LOCATIONS TO SERVE YOU

5645 Blanding Blvd.  
Jacksonville, FL  
771-6330

899 Blanding Blvd.  
Orange Park, FL  
272-1230

11524 San Jose Blvd.  
Mandarin, FL  
262-1965

NET WT. 22 LBS. 8 OZ.

Treats 3,000 Sq. Ft.

Net Wt. 22 Lbs. 18 Oz. 2 3 4

HALL'S  
WEED & FEED  
FOR ST. AUGUSTINE

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS

C A U T I O N

Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

Atrazine can leach through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For best results apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in spring and fall. On newly sprigged St. Augustine, Centipede or Zoysia lawns, apply immediately after planting to prevent weeds from retarding growth. Use herbicides properly.

Do not over-apply. Read directions for use carefully. Water in well. Apply between January 1 and March 31 (Florida and other Gulf states to June 15th).

Apply 7 1/2 lbs. per 1,000 sq. ft. (20 ft. X 50 ft. area). Contents of bag treats 3,000 sq. ft.

Approximately 4 weeks after initial treatment with product, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

**BEST AVAILABLE COPY**

**CONTROLS THESE WEEDS**

Germinating: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings (Dallisgrass, Vaseygrass, Bahiagrass). Sedges (such as Cyperus Compressus), Spurge and Sandburs.

Established and Germinating: Betony, Blackmedic (Yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort (Dollarweed) and Creeping Charlie (Matchweed).

Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bencgrass, or Carpetgrass or where clover is present and desired.

Avoid application to the root zones of trees and shrubs.

Avoid applications on garden plants, flowers and shrubs.

Delay spring planting for one month after application. Do not apply in areas underlaid by roots of desirable trees and shrubs. Clean spreader after use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during the months of July and August.

**SPREADER SETTINGS**

SPREADER	SETTING
Cyclone	4 1/4
Scotts PF-1	4

NOTE: Spreader rate varies with walking speed, check spreader over known area.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.	Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.
	Container Disposal: Do not re-use bag. Discard bag in trash.

**GUARANTEED ANALYSIS**

735-8125

TOTAL NITROGEN (N) .....	21.00%
Ammoniacal Nitrogen .....	1.56%
Water Soluble Organic Nitrogen..	19.44%
AVAILABLE PHOSPHORIC ACID (P <sub>2</sub> O <sub>5</sub> ) .....	4.00%
SOLUBLE POTASH (K <sub>2</sub> O).....	4.01%
Chlorine, not more than .....	2.00%

Derived from Urea, DiAmmonium Phosphate, Triple Superphosphate, Sulphate of Potash.

**BEST AVAILABLE COPY**

494

Statement of Secondary Plant Nutrients:

Iron as Fe ..... 1.50%

Derived from Iron Oxide and Iron Sulphate.

C A U T I O N

Keep out of reach of children. Most fertilizers may stain patios and sidewalks. Remove spillage promptly.

Manufactured by Vigoro Industries, Inc.  
Fairview Heights, IL 62208

