

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH 567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE SEP 27 1984
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
*(Under the Federal Insecticide, Fungicide,
 and Rodenticide Act, as amended)*

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

BEST AVAILABLE COPY

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

[Faint, mostly illegible text follows, likely containing registration details and conditions of use.]

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE SEP 27 1984
---------------------------------	----------------------------

- d. Revise the storage and disposal statement to read:

STORAGE: Store in a dry place. Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. ~~Then dispose of empty bag into application equipment.~~ Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Richard F. Mountford *RFM*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

FRONT PANEL

ACCEPTED
with comments
SEP 27 1984 Rth
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under 557-1966
EPA Reg. No. 15-0-14
TREATS
up to 5,000 sq.ft.

SLOW RELEASE FERTILIZER
PLUS RONSTAR®*
PAR EX®
PROFESSIONAL PRODUCTS

Or Name? ^{H?}
of the
Ronstar
G-label

Provides uniform continuous feeding as it controls weeds in established Bermuda Grass, St. Augustine Grass, Zoysia Grass, Perennial Bluegrass and Perennial Ryegrass.

ACTIVE INGREDIENT

**Oxadiazon (2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)-1,3,4-oxadiazolin-5-one) 0.69%



INERT INGREDIENTS 99.31%
Total 100.00%

**Contains 13.8 pounds of Technical RONSTAR per ton.

For sale to, use and storage by professional turf maintenance personnel only.

KEEP OUT OF REACH OF CHILDREN
WARNING
In case of contact: for eyes, flush with water for 15 minutes; for skin, wash with plenty of soap and water. Get medical attention if irritation persists. See back panel for additional precautionary statements.

*Registered trademark of Rhone-Poulenc, Inc. E.P.A.Reg.No.557- EPA Est.No.557r.

NET WT 50 LBS

REAR PANEL

(1)

NET WT 50 LBS

15-0-14

SLOW RELEASE FERTILIZER

PLUS RONSTAR®

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 15.0%

12.0% Ammoniacal Nitrogen
2.5% Water Insoluble Nitrogen
0.5% Water Soluble Urea Nitrogen

SOLUBLE POTASH (K₂O) 14.0%

Derived from sulfate of ammonia, isobutylidene diurea, muriate of potash and sulfate of potash-magnesia.

Statement of Secondary Nutrients

Total and Water Soluble Magnesium (Mg) 0.50%
Sulfur, combined (S) 14.00%
Iron (Fe) 0.70%
Manganese (Mn) 0.80%

Derived from sulfate of ammonia, sulfate of potash-magnesia, fritted iron and fritted manganese.

Chlorine, not more than 12.00%

PRECAUTIONARY STATEMENTS


HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Harmful if swallowed. Avoid breathing dust. Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Wash contaminated clothing before re-use.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.



DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. Rake area free of leaves.
2. Mow, if necessary, so granules reach soil surface.
3. Measure area to be treated and apply required amount uniformly, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results.
4. Water-in promptly after application.

Par Ex Slow Release Fertilizer plus Ronstar will give pre-emergent control of the following weeds in established Bermuda Grass, St. Augustine Grass, Zoysia Grass, Perennial Bluegrass and Perennial Ryegrass on golf-course fairways, parks and lawns.

Crabgrass	Oxalis
Annual bluegrass	Stinging nettle
Goosegrass	Carpetgrass
Florida Pusley	Pigweed

Amount to Use

Apply prior to weed seed germination.

<u>Weed</u>	<u>When to use</u>	<u>Amount</u>
Goosegrass, Crabgrass	In early spring	6-2/3 lbs. to 10 lbs/1,000 sq. ft.
Annual Bluegrass	In fall or early spring	13-1/3 lbs. per 1,000 sq. ft.
Pigweed, Florida Pusley, Oxalis, Stinging Nettle and Carpetgrass	In early spring	13-1/3 lbs. per 1,000 sq. ft.

PRECAUTIONS.

DO NOT apply to wet turf as injury may result.
 DO NOT apply to Dichondra or Centipedegrass. DO NOT use on putting greens or tees or on turf areas containing bentgrass or red fescue. DO NOT apply to newly seeded areas. Seeding into treated areas should be delayed for 120 days after treatment. Do not apply to turf with heavy thatch accumulation or to turf which has been affected by disease or nematode activity. DO NOT feed treated clippings to livestock. *or allow domestic animals to graze treated areas*
 This product will stain patios and sidewalks. Remove spillage promptly.

Handwritten notes:
 4 1/2 lbs
 270
 .69 lb
 13.04 lbs
 not registered with

STORAGE AND DISPOSAL

STORAGE: - Store in a dry place. Do not contaminate feed and foodstuffs, seeds, fertilizers or other pesticides.

DISPOSAL: - Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

update

Manufactured by Estech, Inc.
Chicago, Illinois 60602

