

FRONT PANEL

10-6-4  
Treats up to  
5,000 sq.ft.

I D E A L 18  
INSECT CONTROL PLUS  
LAWN FERTILIZER



Controls Certain White Grubs, Chinch Bugs,  
Brown Dog Ticks, Cutworms and other insects  
in lawns and turfgrasses

Keep out of reach of children  
**CAUTION**  
STATEMENT OF PRACTICAL TREATMENT  
If on skin: Wash with soap and  
water. If in eyes: Flush with  
plenty of water. See back panel  
for additional precautionary  
statements.

ACTIVE INGREDIENTS

Chlorpyrifos (0,0-diethyl-0-  
(3,5,6-trichloro-2-pyridyl)  
phosphorothioate ..... 0.64%


INERT INGREDIENTS ..... 99.36%

Total ..... 100.00%

U. S. Patent No. 3,244,586  
EPA Reg. No. 557-1980 EPA Est. No. 557-

**ACCEPTED**  
JUL - 6 1990  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 557-1980

NET WT 18 LBS



**BEST AVAILABLE COPY**

REAR PANEL

NET WT. 18 LBS.

Treats up to 5,000 sq. ft.

I D E A L 18

INSECT CONTROL PLUS  
LAWN FERTILIZER

10-6-4

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) .....	10.0Z
10.0Z Ammoniacal nitrogen	
AVAILABLE PHOSPHORIC ACID (P <sub>2</sub> O <sub>5</sub> ) .....	6.0Z
SOLUBLE POTASH (K <sub>2</sub> O) .....	4.0Z

Derived from ammoniated phosphates, ammonium sulfate and muriate of potash.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes or clothing. Avoid breathing of dust. Wash thoroughly after handling. Wash contaminated clothing before re-use. Harmful if swallowed. Do not take internally. Keep children and pets off treated areas until product is washed into the soil and grass is dry.

Note to Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Keep out of lakes, streams or ponds. Do not apply where run-off is likely to occur. This product is toxic to fish, birds and other wildlife. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

**BEST AVAILABLE COPY**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

LAWN AND TURF AREAS: To control Chinchbugs, Sod Webworms, Ants, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Grasshoppers and Sowbugs (Pillbugs), apply the contents of this bag uniformly over 5,000 sq. ft. using the spreader setting shown below, followed IMMEDIATELY by thorough watering to carry the active ingredient into the zone of insect infestation.

To control certain white grubs (larvae of European Chafer and Japanese Beetle), apply the contents of this bag uniformly over 2,500 sq. ft. using the spreader setting shown below, followed IMMEDIATELY by thorough watering. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

Repeat as necessary if re-infestation occurs.

SPREADER SETTINGS

SPREADER	Setting	
	18 lbs/5,000 sq.ft.	18 lbs/2,500 sq. ft.
Cyclone	4	4 1/2
Sears	4 3/4	7
Scotts PF-1	5	6 3/4
Precision 20C	3 3/4	6 1/2

NOTE: Spreader rate varies with walking speed and condition of spreader. Check spreader over measured area and adjust, if necessary, to apply correct amount. Remove spillage promptly.

Approximately 4 weeks after initial treatment with product, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.	Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Container Disposal: Do not reuse bag. Discard bag in trash.
---	--

Manufactured by Vigoro Industries, Inc.  
Fairview Heights, IL 62208

**BEST AVAILABLE COPY**