

PM 25

557-1983

1 of 4

Treats 10,000 sq. ft.

PARK RIDGE WEED & FEED
FOR ST. AUGUSTINE GRASSES
20-2-10

For exclusive use on St. Augustine, Centipede and Zoysia Grasses.

Controls germinating and certain established weeds including Dollar Weed.

Fortified with 1% iron and other micronutrients.

KEEP OUT OF REACH OF CHILDREN

CAUTION


STATEMENT OF PRACTICAL TREATMENT

In case of contact, for skin, wash with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention if irritation persists. See back panel for additional precautionary statements.

ACTIVE INGREDIENT:

*Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine	0.86%
*Related compounds05%
INERT INGREDIENTS	99.09%
TOTAL	100.00%

*Contains 18.2 lbs. of atrazine and related compounds per ton



EPA Reg. No. 557-1983 EPA Est. No. 557-FI-11

NET WT. 50 LBS.

REAR PANEL

Treats 10,000 sq. ft.

NET WT. 50 LBS.

PARK RIDGE WEED & FEED

FOR ST. AUGUSTINE GRASS S

20-2-10

GUARANTEED ANALYSIS

TOTAL NITROGEN (N).....	20.0%
0.90% Ammoniacal Nitrogen	
*16.50% Urea Nitrogen	
2.60% Water Insoluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅).....	2.0%
SOLUBLE POTASH (K ₂ O).....	10.0%

Derived from ammoniated phosphate, isobutylidene diurea, sulphur coated urea, activated sludge, muriate of potash and sulphate of potash-magnesia.

Secondary Nutrients

Magnesium (Mg)	1.0%
1.0% Water Soluble Magnesium (Mg)	
Sulphur, combined (S)	2.0%
Sulphur, free (S)	2.0%
Iron (Fe).....	1.0%
Manganese (Mn).....	1.0%

Derived from sulphate of potash-magnesia, sulphur coated urea, iron oxide, iron sulphate, manganous oxide and manganese sulphate.

*Includes 4.0% coated slow release urea nitrogen from sulphur coated urea.

Chlorine, not more than	6.5%
-------------------------------	------

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS & DOMESTIC ANIMALS

C A U T I O N

Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Keep out of reach of children. Do not take internally.

BEST AVAILABLE COPY

ENVIRONMENTAL HAZARD

Atrazine can leach through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not contaminate water when disposing of equipment washwaters. Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass or Carpetgrass or where clover is present and desired.

Avoid application to the root zones of trees and shrubs. Avoid applications on garden plants, flowers and shrubs. Delay spring planting for one month after application. Clean spreader after each use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during months of July, August and September

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply between January 1 and March 31 (Florida and other southern states to June 15th) at the rate of 50 lbs. on 10,000 sq. ft.

CONTROLS SUCH WEEDS AS:

GERMINATING: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings... (Dallisgrass, Vaseygrass, Bahiagrass), Sedges (such as Cyperus Compressus), and Spurge.

ESTABLISHED AND GERMINATING: Betony, Blackmedic (yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort (Dollarweed) and Creeping Charlie (Matchweed).

For best results, apply before weeds come up (pre-emergent). Established weeds are more difficult to control and may require an application in Spring and Fall. On newly sprigged St. Augustine, Centipede, or Zoysia lawns, apply immediately after planting to prevent weeds from retarding growth. Water in well.

Approximately 4 weeks after initial treatment with product, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

BEST AVAILABLE COPY

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not re-use bag. Discard bag in trash.

SPREADER SETTINGS

SPREADER	SETTING
Cyclone	5 1/2
Scott R-8	M
Lasco	J

NOTE: Spreader setting varies with terrain, equipment and walking speed. To insure proper coverage, calibrate before use by checking over measured area.

CAUTION: Close spreader openings when loading, turning or when stopped. This product will stain patios, sidewalks and swimming pools. Sweep spillage promptly.

Park Ridge® is a registered trademark of Vigoro Industries, Inc.

Manufactured by
Vigoro Industries, Inc.
Fairview Heights, IL 62208

BEST AVAILABLE COPY