

PM-25

Reg # 557-1984

194

Environmental Protection Agency
Washington, DC 20460

Registration

Standardization

Other

JAN - 8 1997

Mr. Michael J. Archer
IMC Vigoro
P.O. Box 512
Winter Haven, FL 33882

Dear Mr. Archer:

Subject: Minor Changes
IMC Vigoro
EPA Reg. No. 557-1984
Your submission dated Dec. 12, 1996

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable and a stamped copy is enclosed for your records.

Sincerely yours,

Philip V. Errico
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

294

FRONT PANEL

PARK RIDGE 20 WEED & FEED

TREATS 5,000 SQ. FT.

10-4-6

For exclusive use on St. Augustine, Centipede and Zoysia Grasses.
Controls germinating and certain established weeds including Dollar Weed.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, for skin, wash with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention if irritation persists.
See back panel for additional precautionary statements.

ACTIVE INGREDIENT:

*Atrazine: 2-chloro-4-ethylamino-6-isopropylamine-s-atrazine.....	1.00%
*Related compounds.....	0.05%
INERT INGREDIENTS:.....	98.95%
TOTAL.....	100.00%

*Contains 21.0 lbs. of atrazine and related compounds per ton.

EPA Reg. No. 557-1984
EPA Est. no. 557-FL-1

NET WEIGHT 20 LBS.

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated

J. - 8 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
557-1984

96 DEC 18 AM 1:56

RECD 12/18/96 12/18/96

REAR PANEL

PARK RIDGE 20 WEED & FEED FOR ST. AUGUSTINE GRASSES

10-4-6
GUARANTEED ANALYSIS

Total Nitrogen (N).....	10.00%
10.00% Ammonical Nitrogen	
Available Phosphorus (P_2O_5).....	4.00%
Soluble Potash (K_2O).....	6.00%
Derived from ammoniated phosphate, sulfate of ammonia and muriate of potash.	
Chlorine, not more than.....	6.00%

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Keep out of reach of children. Do not take internally.

ENVIRONMENTAL HAZARDS

Atrazine can leach through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well -drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Do not use north of North Carolina. Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass or Carpetgrass or where clover is present and desires. Avoid application to the root zones of trees and shrubs. Avoid application on garden plants, flowers and shrubs. Delay spring planting for one month after application. Clean spreader after each use.

Apply only in Winter, Spring and Fall. This product is not recommended for use during months of July, August and September.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply between January 1 and March 31 (Florida and other southern states to June 15) at the rate of 20 lbs. on 5,000 sq. ft.

CONTROLS SUCH WEEDS AS:

GERMINATING: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings (Dallisgrass, Vaseygrass, Bahiagrass), Sedges (such as Cyperus Compressus), and Spurge.

ESTABLISHED AND GERMINATING: Betony, Blackmedic (yellow trefoil), Burclover, Chickweed, Dandelion, Oxalis, Florida Pursley, Pennywort (Dollarweed) and Creeping Charlie (Matchweed).

494

For best results, apply before weeds come up (Pre-emergent). Established weeds are more difficult to control and may require an application in Spring and Fall. On newly sprigged St. Augustine, Centipede or Zoysia lawns, apply immediately after planting to prevent weeds from retarding growth. Water in well.

For homeowner use only.

Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds food or feed. Return spillage to original container prior to disposal.
PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.
CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash

PRECAUTIONS:

Do not use north of North Carolina.
Do not use on lawns of Bahiagrass, Bermudagrass, Dichondra, Bentgrass or Carpetgrass or where clover is present and desired.
Avoid application to the root zones of trees and shrubs.
Avoid application on garden plants, flowers and shrubs.
Delay spring planting for one month after application.
Apply only in Winter, Spring and Fall.
This product is not recommended for use during months of July, August and September.

SPREADER SETTINGS

SPREADER	SETTING
Central 40 Drop	5 1/2
Central Broadcast	4 1/4
Cyclone	4 1/4
Greenview	7 1/2
Ortho 3000	7 1/2
Scotts Drop	6 1/2
Sears Drop	5 1/4
Sears Broadcast	4 1/4

(Note to regulators: Spreader listing and Setting may be changed to reflect market changes.)

NOTE: Spreader setting varies with terrain, equipment and walking speed. To insure proper coverage, calibrate before use by checking over measured area.

CAUTION: Close spreader openings when loading, turning or when stopped. This product will stain patios, sidewalks and swimming pools. Sweep spillage promptly.

Park Ridge® is a registered trademark of Vigoro Industries, Inc.
Manufactured by IMC Vigoro, PO Box 512, Winter Haven FL 33882