

FRONT PANEL

TREATS 5000 SQ. FT.

VIGORO®

DEEP GREEN® WEED & FEED

FOR BAHIA, BERMUDA AND ZOYSIA GRASSES

KEEP OUT OF REACH OF CHILDREN

CAUTION

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention. See back panel for additional precautionary statements.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.408%
+ **Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	0.117%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.050%
INERT INGREDIENTS	99.425%
TOTAL	100.000%


Used at 25 lbs. on 5000 sq. ft. equivalent to:

- *MCPA Acid - 0.338% - 0.74 lbs. per acre or 6.76 lbs. per ton.
- + **2,4-D Acid - 0.097% - 0.21 lbs. per acre or 1.94 lbs. per ton.
- ***Dicamba Acid - 0.041% - 0.09 lbs. per acre or 0.82 lbs. per ton.
- +Isomer specific by AOAC Method 6D01.5

J. S. Patent No. 3,284,186
EPA Reg. No. 557-

EPA Est. No. 557-FL-1

NET WT. 25 LBS.



BEST AVAILABLE COPY

REAR PANEL

NET WT. 25 LBS.

Treats 5000 sq. ft.

VIGORO[®]

DEEP GREEN[®] WEED & FEED

FOR BAHIA, BERMUDA AND ZOYSIA GRASSES

19-2-6

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	+ 19.00%
9.00% Ammoniacal Nitrogen	
8.40% Urea Nitrogen	
1.60% Water Insoluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	2.00%
SOLUBLE POTASH (K ₂ O)	6.00%
Derived from sulphate of ammonia, isobutylidene diurea, sulphur coated urea, diammonium phosphate, muriate of potash and sulphate of potash-magnesia.	
Secondary Nutrients:	
Total Magnesium (Mg)	1.00%
1.00% Water Soluble Magnesium (Mg)	
Iron (Fe)	1.50%
Total Manganese (Mn)	0.30%
Derived from sulphate of potash-magnesia, iron oxide and manganous oxide.	
*This product contains 7.70 units of slow release nitrogen from isobutylidene diurea and sulphur coated urea.	
Chlorine, not more than	4.00%

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Keep out of reach of children. Wash thoroughly after handling. Harmful if swallowed. Do not re-enter or permit others to re-enter treated areas until dusts have settled.

Protective Clothing: Applicators must wear long pants, long-sleeved shirts, shoes and gloves.

BEST AVAILABLE COPY

REAR PANEL

ENVIRONMENTAL HAZARD

This product is toxic to aquatic invertebrates. Drift or run-off may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. Do not contaminate irrigational ditches or water used for domestic purposes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. Mow lawn to normal height 1 or 2 days before application.
2. Water lawn thoroughly 1 day before application to ensure adequate soil moisture. Do not apply to wet grass.
3. Apply contents of this bag to 5,000 sq. ft. using the suggested spreader setting. DO NOT OVERAPPLY. For best results, apply when weeds are young and actively growing and the average temperature is over 70° F.
4. Do not mow or water lawn again until 2 days after application.
5. Approximately 4 weeks after weed kill is evident, an application of 16% or higher nitrogen fertilizer should be made at a rate of 6 lbs. per 1,000 sq. ft. and watered in well.

This product may be used on Bahia, Bermuda, Zoysia, Fescue, Kentucky bluegrass, St. Augustine, Centipede and Bentgrass lawns. If adverse low moisture conditions exist, weeds are in a poor state of growth, or area is heavily overrun with weeds, retreatment may be necessary. Make a second application in approximately 30 days.

PRECAUTIONS: Do not apply to newly seeded lawns. Do not apply to dichondra, lippia or other ground cover. Avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly because temporary foliage discoloration may occur. Do not use where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables or ornamental shrubbery. Apply when air is calm. Do not apply to areas underlaid by roots of desirable trees and shrubs.

<p>Storage: Store in original container in safe, dry place. Do not contaminate other pesticides, fertilizer, seeds, food or feed. Return spillage to original container prior to disposal. Keep bag tightly closed. Do not store near growing plants.</p>	<p>Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.</p> <p>Container Disposal: Do not re-use bag. Discard bag in trash.</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

PRO

TON

REAR PANEL Cont.

CONTROLS THESE WEEDS

Florida Pusley, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida Spurge, Wild Onion, Wild Lettuce and most other broadleaf weeds.

SPREADER SETTING

SPREADER	SETTING	SPREADER	SETTING
Vigoro Rotary	5	Precision 20C.	6 1/4
Cyclone	5 1/2	Scotts PF-1	6 1/2
Sears.	6 1/2	Scotts R-8	M

Note: Spreader rate varies with walking speed, check spreader over known area. Close spreader openings when loading, turning or when stopped. Hand spreading is not recommended.

Clean spreader after each use.

Manufactured by Vigoro Industries, Inc., Fairview Heights, Il 62208. Vigoro® and Deep Green® are registered trademarks of Vigoro Industries, Inc.