

Reg # 557-1990

PM-23

19 15

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (11-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.

557-1990

DATE OF ISSUANCE

24 SEP 1992

TERM OF ISSUANCE

CONDITIONAL, EX. 1000: JAN. 1, 1995

NAME OF PESTICIDE PRODUCT

100% SOLUBLE PHOSPHATE FERTILIZER
FERTILIZER WITH NITROGEN
FERTILIZER

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Widoro Industries, Inc.
P.O. Box 4139
Fairview Heights, IL 62206

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with section 5(a)(7)(C) of the Act.

- 1. The label for this product shall be identical to the label for the product registered under registration number 557-1990.
- 2. The label for this product shall be identical to the label for the product registered under registration number 557-1990.

The registrant shall, within 90 days of the date of this notice, submit to the Administrator a copy of the label for this product for review and approval.

The registrant shall, within 90 days of the date of this notice, submit to the Administrator a copy of the label for this product for review and approval.

Scanned by Miller
Product Manager (23)
Fungicide-Rodenticide Branch
Registration Division (11-767)

ENCLOSURE

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

[Handwritten Signature]

DATE

7/22/92

2915

**PAR EX® SLOW RELEASE FERTILIZER
with BARRICADE® HERBICIDE**

**For selective preemergence control of grass and broadleaf weeds in
established turf grasses and landscape ornamentals.**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**See the following section, Precautionary Statements, Hazards to Humans
and Domestic Animals, for additional information.**

ACTIVE INGREDIENT:

Prodiamine [N³,N³-Di-n-propyl-2,4 dinitro-6-
(trifluoromethyl)-m-phenylenediamine]..... 0.345%

INERT INGREDIENTS:..... 99.655%

EPA Reg. No. 557- R00N

EPA Est. No. 557-FL-1

BARRICADE® is a registered
trademark of Sandoz, Ltd.

NET WT. 50 LBS.

24 SEP 1992

557-1990

3915

NET WT. 50 LBS.

PAR EX® SLOW RELEASE FERTILIZER
with BARRICADE® HERBICIDE

32-4-6

GUARANTEED ANALYSIS

Total Nitrogen (N).....	32.0%
1.5% Ammoniacal Nitrogen	
22.5% Urea Nitrogen	
8.0% Water Insoluble Nitrogen	
- Available Phosphoric Acid (P ₂ O ₅).....	4.0%
Soluble Potash (K ₂ O).....	6.0%
Derived from ammonium phosphate, urea, isobutylidene diurea, ureaform and muriate of potash.	

41 2 15

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

Statements of Practical Treatment

If in eyes: Flush thoroughly with water for several minutes. Get medical attention if irritation persists.

If on skin: Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists.

If inhaled: Remove victim to fresh air.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

5215

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in turf grasses and landscapes. This product controls susceptible weeds by inhibiting weed seeds germination and root development. For most effective weed control this product must be incorporated by 1/2 inch of rainfall, irrigation or shallow mechanical incorporation. Incorporation should take place as soon as possible after application and should not be delayed past 14 days from the date of application.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

APPLICATION DIRECTIONS


Apply uniformly through suitable calibrated commercial application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established grass turf. The species of turf grass determines the maximum amount of material that may be applied during a year.

Most effective weed control in turf grasses will be obtained when this product is activated by a least 1/2 inch of rainfall or irrigation within 14 days after application and prior to weed germination. See the map below for approximate crabgrass germination dates.

CRABGRASS SEED GERMINATION DATES



Approximate Dates

BEST AVAILABLE COPY

-  After May 30
-  After May 10
-  After April 20
-  After March 20
-  January 1 to March 20

Foxtails (yellow, green) tend to germinate before crabgrass; therefore, if targeting foxtails, apply this product three (3) weeks before the corresponding crabgrass germination date.

6915

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE

Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or; puncture and dispose of in a sanitary landfill or at an incineration facility, or; if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

Paper and plastic containers: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or at an incineration facility, or; if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

Fiber Drums: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residues into application equipment. Dispose of empty liner in a sanitary landfill or at an incineration facility if allowed by state and local authorities. If the drum cannot be reused, dispose of in a sanitary landfill or by incineration if allowed by state and local authorities.

BEST AVAILABLE COPY

7 9 15

MAXIMUM ANNUAL RATES

This product should not be applied above a maximum annual rate of 435 lbs. per acre or 10 lbs./1000 square feet.

TURF SPECIES AND RATES

For season long control this product may be applied as a single application, or if needed, followed by a second application for increased length of control. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rate within the listed range for longer control. Longer control is desirable in areas where crabgrass germinates between January 1 and April 20 (See Crabgrass Seed Germination Data Map).

Select maximum use rate based on turf grass species. All applications must be made prior to germination of the target weed seeds.

The rates listed in the table below are for weed control of approximately 2-6 months.

TURF SPECIES AND RATES TABLE

TURF SPECIES: =====	Minimum & Maximum Rate of Product Per Calendar Year	
	Lbs./Acre	Lbs./1000 Sq. Ft.
Bermudagrass Centipedegrass Seashore Paspalum Tall Fescue Zoysia -----	93 - 435	2 - 10
Bluegrass Perennial Ryegrass -----	93 - 290	2 - 6 2/3
Creeping Red Fescue St. Augustinegrass -----	93 - 214	2 - 5
Creeping Bentgrass min. 1/2 inch height -----	93 - 191	2 - 4 2/5
Bermudagrass newly sprigged or plugged ¹	93 - 151	2 - 3 1/2

¹ Suppression only of Foxtail, Goosegrass and Rescuegrasses due to reduced product rates.

BEST AVAILABLE COPY

- Do not apply more than 217.5 lbs. of this product per application.
- Do not make more than 2 applications per calendar year.
- Do not apply more than 435 lbs. of this product per calendar year.
- If making two applications, wait at least 60 days after the first application before making a second application.
- Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type.

WEEDS CONTROLLED

When used at the recommended rates (see Turf Species and Rates Table) this product will control:

- | | |
|---------------------------|----------------------|
| Annual Bluegrass | Knotweed |
| Common Chickweed (1) | Oxalis |
| Crabgrass (Large, Smooth) | Prostrate Spurge |
| Foxtails (Yellow, Green) | Rescuegrass (3) |
| Goosegrass (2) | Shepherd's purse (1) |
| Henbit | |

- (1) Fall application.
- (2) For the best control of goosegrass, apply 217.5 lbs/acre. Sixty (60) days later make a second application with 72.5 lbs. to 217.5 lbs./acre of product (see information in Turf Species and Rates section above).
- (3) Suppression only.

The following use precautions apply to the use of this product in turf grasses:

Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to FALL SEEDED TURF until the following spring and only after the second mowing.

Do not apply to SPRING SEEDED TURF or newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Do not apply this product to tees or greens, or to areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (Poa annua) are desirable species.

OVERSEEDING/RESEEDING

This product will inhibit the germination of turf species if overseeded too soon after application.

The interval between application and overseeding is dependent on the application rate, soil temperature, and soil moisture. A shorter interval is required if growing conditions are warm and wet, while a longer interval is needed if conditions are cool and dry.

Use the Overseeding/Reseeding Intervals Table below as a guide when choosing the appropriate overseeding/reseeding interval for each situation.

Overseeding/Reseeding Intervals Table:

Rate Lbs./Acre	Overseeding/Reseeding Interval (Months) Based Upon Seeding Method and Predominant Soil Conditions for the Season			
	Broadcast Seeding		Drill Seeding	
	Wet & Warm	Cool & Dry	Wet & Warm	Cool & Dry
93-145	4	6	3	4
146-290	8	12	6	8
291-435	12	18	9	12

This product will not harm most established ornamental trees, shrubs, and flowers when applied to surrounding areas in accordance to the directions given in the Turf Species and Rates section. The following species have shown tolerance to this product.

<u>Scientific name</u>	<u>Common name</u>
<i>Acer palmatum</i>	Japanese Maple
<i>A. platanoides</i>	Norway Maple
<i>Actinidia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Barberis gladwynensis</i>	
<i>B. julianae</i>	Wintergreen Barberry
<i>B. mentorensis</i>	Mentor Barberry
<i>B. thunbergii</i>	Japanese Barberry
<i>B. verruculosa</i>	Warty Barberry
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisoides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Cleyera japonica</i>	Cleyera
<i>Cornus florida</i>	Flowering Dogwood
<i>C. stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>C. buxifolius</i>	
<i>C. dammeri</i>	Bearberry Cotoneaster
<i>C. microphyllus</i>	Rockspray Cotoneaster
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delasperma alba</i>	White Trailing Ice Plant
<i>Dodonea viscosa</i>	Hop Bush
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper Euonymus

*Ornamental species only. Do not use on food producing trees or vines.

BEST AVAILABLE COPY

Scientific name**Common name**

E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
F. viridissima	Greenstem Forsythia
Gardenia jasminoides	Gardenia, Cape-Jasmine
Hedera helix	English Ivy
Hibiscus rosa-sinensis	Chinese Hibiscus
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. pernyi	
Jasminium nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper
J. conferta	Shore Juniper
J. horizontalis	Creeping Juniper
Juglans sp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crepe Myrtle
Ligustrum amurense	Amur Privet
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet
Liriope muscari	Big Blue Lily Turf
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Maleophora luteola	Ice Plant
Malus sp.	Crabapple*
Nandina domestica	Heavenly Bamboo
Olea europa	Olive*
Ophiopogon japonicus	Mondo Grass
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
Persea americana	Avocado*
Photinia fraseri	Fraser's Photinia
Picea abies	Norway Spruce
Pieris japonica	Lily-of-the-Valley Shrub

*Ornamental species only. Do not use on food producing trees or vines.

Scientific name

Common name

<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Black Pine
<i>P. radiata</i>	Monterey Pine
<i>P. thunbergiana</i>	Japanese Black Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>Pistachio</i> sp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Mock Orange
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> sp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pyracantha coccinea</i>	Firethorn
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus</i> sp.	Bradford Pear sp.
<i>Quercus rubra</i>	Red Oak
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Rhododendron</i> (including <i>Azalea</i>)	'Coral Bells'
	'Formosa'
	'Hino-crimson'
	'PJM'
	'Roseum Elegans'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary
<i>Rumohra adiantiformis</i>	Leatherleaf Fern
<i>Santolina virens</i>	
<i>Sedum album</i>	Stonecrop
<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum

*Ornamental species only. Do not use on food producing trees or vines.

Scientific nameCommon name

V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush
V. wrightii	
Vinca minor	Dwarf Periwinkle
Vitis sp.	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet
Yucca filamentosa	Yucca
	Adam's Needle

*Ornamental species only. Do not use on food producing trees or vines.

11 9 15

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of manufacturer or seller. All such risks shall be assumed by buyer or user.

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. Manufacturer makes no other express or implied warranty of fitness or of merchantability or any other express or implied warranty. In no event shall manufacturer or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product.

Manufacturer and seller offer this product, and buyer and user accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.

Manufactured by Vigoro Industries, Inc., Winter Haven, FL 33882.
Par Ex[®] is a registered trademark of Vigoro Industries, Inc.

15 7 15

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of manufacturer or seller. All such risks shall be assumed by buyer or user.

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. Manufacturer makes no other express or implied warranty of fitness or of merchantability or any other express or implied warranty. In no event shall manufacturer or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product.

Manufacturer and seller offer this product, and buyer and user accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.

Manufactured by Vigoro Industries, Inc., Winter Haven, FL 33882.
Par Ex® is a registered trademark of Vigoro Industries, Inc.

