

SOLD TO VETERINARIANS ONLY

CODE 0157

ComBot™

Brand of Trichlorfon

EQUINE ANTHELMINTIC

ComBot EQUINE ANTHELMINTIC IS A READY-TO-USE PREPARATION THAT REQUIRES NO FURTHER DILUTION BEFORE USE IN HORSES AND PONIES.

INDICATIONS: For removal and control of Bots (*Gastrophilus* Spp.), Ascarids (*Parascaris equorum*), and Pinworms (*Oxyuris equi*) in horses and ponies.

DIRECTIONS FOR USE AND DOSAGE: Administer by stomach tube at the rate of one half (1/2) fluid ounce of ComBot per 100 pounds of body weight. This will provide a dosage of 18.2 mg of active ingredient per pound of body weight which will effectively control the parasites indicated. Do not fast animals prior to treatment.

Use in Conjunction with Other Anthelmintics: In clinical trials, ComBot at the recommended dosage has been used successfully with the recommended rates of mebendazole, thiabendazole, phenothiazine and piperazine. If ComBot is used with these compounds to increase the spectrum of anthelmintic activity, administer simultaneously but do not mix ComBot with any of these products and store for future use.

CONTRAINDICATIONS: Use only as directed and do not exceed recommended dose. Do not treat colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses or ponies. Horses severely debilitated, suffering from diarrhea or severe constipation, infectious disease, toxemia or colic should not be treated until such conditions are corrected with proper therapy. ComBot is a cholinesterase inhibitor. Do not use this product in horses simultaneously or within 10 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

ComBot is a TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

For removal and control of Bots (*Gastrophilus spp.*),
Ascarids (*Parascaris equorum*), and Pinworms (*Oxyuris equi*)

GUARANTEE

ACTIVE INGREDIENT:

Trichlorfon (Dimethyl [2,2,2-trichloro-1-hydroxyethyl] phosphonate) ----- 12.3% w/v

INERT INGREDIENTS: ----- 87.7% w/v

100.0% w/v

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

WARNING

Not for use on horses intended for food
KEEP OUT OF REACH OF CHILDREN
Do not store product above 86°F
Protect from sunlight

NET CONTENTS — 1 GALLON

SIDE EFFECTS: Occasionally slight transient after administering ComBot. Fasting prior increase the incidence of these side effects.

NOTICE TO VETERINARIAN: Symptoms of severe colic, muscle tremors and ataxia. At

WARNING: Not for use in horses intended for

PRECAUTION: Harmful if swallowed. Avoid soap and water after use. If swallowed by tablespoonfuls of salt in a glass of water.

TO PHYSICIAN: This material is a cholinesterase inhibitor. 2-PAM is also antidotal and may be administered as supportive treatment. Do not walk patient

EPA Reg. No. 569-68

EPA Est. No. 3125-MO-1

Haver-Lockhart
LABORATORIES

CODE 0157

ComBotTM

Brand of Trichlorfon
EQUINE ANTHELMINTIC

SOLD TO VETERINARIANS ONLY

For removal and control of Bots (*Gastrophilus spp.*),
Ascarids (*Parascaris equorum*), and Pinworms (*Oxyuris equi*)

GUARANTEE

ACTIVE INGREDIENT:

Trichlorfon (Dimethyl [2,2,2-trichloro-1
-hydroxyethyl] phosphonate) ----- 12.3% w/v

INERT INGREDIENTS: ----- 87.7% w/v
100.0% w/v

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the
order of a licensed veterinarian.

WARNING

Not for use on horses intended for food

KEEP OUT OF REACH OF CHILDREN

Do not store product above 86°F
Protect from sunlight

NET CONTENTS — 1 GALLON

SIDE EFFECTS: Occasionally slight transient diarrhea and/or colic may occur soon
after administering ComBot. Fasting prior to the administration of ComBot will
increase the incidence of these side effects.

NOTICE TO VETERINARIAN: Symptoms of overdose in horses are profuse diarrhea,
severe colic, muscle tremors and ataxia. Atropine is antidotal.

WARNING: Not for use in horses intended for food. Keep out of reach of children.


PRECAUTION: Harmful if swallowed. Avoid contact with the skin. Wash hands with
soap and water after use. If swallowed by humans, induce vomiting by giving 2
tablespoonfuls of salt in a glass of water. Call physician immediately.

TO PHYSICIAN: This material is a cholinesterase inhibitor. Atropine is antidotal.
2-PAM is also antidotal and may be administered in conjunction with atropine. Give
supportive treatment. Do not walk patient nor give morphine.

EPA Reg. No. 569-68

EPA Est. No. 3125-MO-1

Haver-Lockhart
LABORATORIES



12-1-68 569-68