



New improved

# ROCKLAND PROlin<sup>®</sup> Concentrate

Mix Your Own Rat and Mice Bait



ACTIVE INGREDIENTS:	
3(a-acetonylbenzyl)-4-hydroxycoumarin (warfarin)	0.5%
N <sup>2</sup> -2-quinazolylsulfanilamide (sulfaquinoxaline)	0.5%
INERT INGREDIENTS:	99.0%
	100.0%

Net Weight 4 Oz. Makes 5 lbs. of Bait

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
SEE SIDE PANEL FOR COMPLETE CAUTION STATEMENT.

\*Trade mark of the Wisconsin Alumni Research Foundation

Manufactured for

Rockland Chemical Co., Inc  
PASSAIC AVENUE WEST CALDWELL, N.J.

### CAUTION:

Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin. It hereby increases effectiveness against hard-to-kill rats and mice and gives better and more thorough control.

U.S.D.A. 572-31

### DIRECTIONS FOR MANUFACTURING RAT OR MOUSE BAIT:

Mix Prolin<sup>™</sup> Concentrate thoroughly with cracked or ground grains, dog food, or other edible inert baiting materials. Use 1 part Prolin Concentrate to 19 parts by weight of inert bait.

### DIRECTIONS FOR USE OF RAT BAIT:

Place 4 to 6 ounces of bait in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

### DIRECTIONS FOR USE OF MOUSE BAIT:

Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 12 days.

ACCEPTED  
APR 10, 1967  
UNDER THE FEDERAL  
PURCHASE AND  
RENTAL ACT  
FOR SUPPLYING  
ED UNDER  
572-31

Remnant 5

12

**RC** **ROCKLAND**  
**New improved PROLIN Concentrate**  
**Mix Your Own Rat and Mice Bait**

**ACTIVE INGREDIENTS:**  
 3-(p-aminophenyl)-2,4-dihydroxyquinoline (warfarin) 8 1/2%  
 N-(2-ethylhexyl)-2,4-dinitrophenylamine (difenacoum) 91 1/2%

**INERT INGREDIENTS:** 100%

Net Weight 4 Oz. Makes 5 lbs. of Bait

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
 SEE SIDE PANEL FOR COMPLETE CAUTION STATEMENT.  
 \*Trade mark of the Warfarin & Coumatral Synthesis Corporation

Manufactured for  
**Rockland Chemical Co. Inc.**  
 2011 E. 10th St. - St. Paul, Minn.



**CAUTION:**  
 Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, introduce oral administration of vitamin K combined with blood transfusions as indicated in the case of hemorrhage caused by overdose of dihydrocoumatral.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin. It thereby increases effectiveness against hard-to-kill rats and mice and gives better and more thorough control.

U.S.D.A. 572-31

**DIRECTIONS FOR MANUFACTURING RAT OR MOUSE BAIT:**  
 Mix Prolin Concentrate thoroughly with treated or ground grain, dog feed, or other edible inert baiting materials. Use 1 part Prolin Concentrate to 19 parts by weight of inert bait.

**DIRECTIONS FOR USE OF RAT BAIT:**  
 Place 4 to 6 ounces of bait in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

**DIRECTIONS FOR USE OF MOUSE BAIT:**  
 Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 12 days.

UNDER THE PROVISIONS OF THE FEDERAL FOOD, DRUG AND COSMETIC ACT, THIS PRODUCT IS SUBJECT TO INSPECTION BY THE U.S. DEPARTMENT OF HEALTH, EDUCATION AND WELFARE.

**ACCEPTED**  
 Apr. 10, 1967

*Comments*

13

16