

**DIRECTIONS**

**FOR FLIES**

For the reduction of flies in and around poultry buildings and yards, and around homes, use 2 1/4 gallons in 100 gallons of water, or 1 pint in 5 gallons of water. Apply sufficient spray to thoroughly wet surfaces on which flies are seen to collect, such as walls, ceilings, manure piles, garbage cans, fences. Repeat applications as necessary.

**CAUTION**

Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid repeated or prolonged contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Do not use in milk processing rooms or dwellings.

NOTE TO PHYSICIAN: Malathion, upon repeated, prolonged or careless use, may cause cholinesterase inhibition. Atropine is antidotal.


**FLAMMABLE**

Keep container closed and away from heat and open flame.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ACCEPTED  
J 10 1952

# ROCKLAND MALATHION 50% EMULSIFIABLE CONCENTRATE



For Lice & Red Mites in Poultry, Apply to Roosts & Houses

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
SEE SIDE PANEL FOR COMPLETE CAUTION STATEMENT.

<b>ACTIVE INGREDIENTS:</b>		
*Malathion	-----	50 %
Xylolj	-----	40.8%
<b>INERT INGREDIENTS</b>		9.2%
*O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.		

**Rockland Chemical Co., Inc.**  
PASSAIC AVENUE, WEST CALDWELL, N. J.

ACCEPTED WITH COMMENTS


For control  
Mix Rockla  
mixing tabl  
or a house  
Force spray  
application:  
To make a  
oil or wate  
To make 1%  
to 1 part R

ROOST SPR  
rate of 1 p  
HOUSE OR  
surfaces th  
Apply a 2%  
and nests,

ROOST PA  
proximately  
lice and mi  
HOUSE OR  
point of ru  
spraying. V  
a 2% spray.

# ROCKLAND MALATHION 50%

## EMULSIFIABLE CONCENTRATE



For Lice & Red Mites in Poultry, Apply to Roosts & Houses

**CAUTION:** KEEP OUT OF REACH OF CHILDREN.  
SEE SIDE PANEL FOR COMPLETE CAUTION STATEMENT.

**ACTIVE INGREDIENTS:**

*Malathion	50 %
Xylol	40.8%
<b>INERT INGREDIENTS</b>	9.2%
*O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.	

**Rockland Chemical Co., Inc.**  
PASSAIC AVENUE, WEST CALDWELL, N. J.

**DIRECTIONS**

For use as a roost spray or paint, mix 2 parts of Rockland Malathion 50% concentrate with 98 parts of water. For use as a house or premise spray, mix 2 parts of Rockland Malathion 50% concentrate with 98 parts of water. For use as a roost spray, apply directly to roosts, dropping boards and nests, wetting surface to a point of run off. For use as a house or premise spray, clean all surfaces thoroughly and remove old nesting material. Apply a 2% spray directly to roosts, dropping boards and nests, wetting surface to a point of run off. To make a 2% spray, add 2 parts of Rockland Malathion 50% concentrate to 98 parts of water. To make 1% spray add 49 parts of water to 1 part Rockland Malathion 50%.

**FOR LICE**

**ROOST SPRAY OR PAINT:** Apply a 2% mixture at the rate of 1 pint per 50 lineal feet.

**HOUSE OR PREMISE SPRAY:** Before spraying, clean all surfaces thoroughly and remove old nesting material. Apply a 2% spray directly to roosts, dropping boards and nests, wetting surface to a point of run off.

**MITES**

**Red Mite or Roost Mite**

**ROOST PAINT OR SPRAY:** Apply a 1% solution approximately 1 pint to 150 lineal feet. To control both lice and mites use a 2% solution.

**HOUSE OR PREMISE SPRAY:** Spray a 1% solution to point of run off, being certain to clean up before spraying. When controlling both lice and mites use a 2% spray.

U.S.D.A. Reg. No. 572-47