

ROCKLAND ROTENONE 1% DUST is recommended for control of pests as shown below:

[572-53]

PEST

Cabbage Loopers	Cabbage, Broccoli, Cauliflower, Lettuce
Diamond Back Moth	Beans
Mexican Bean Beetle	Potatoes, Tomatoes
Colorado Potato Beetle	Asparagus
Asparagus Beetle	Tomatoes, Peppers, Eggplants
Tomato Hornworm	Cucumber, Squash, Melons
Cucumber Beetle	Potatoes, Peppers, Eggplants, Sweet Potatoes, Strawberries, Grapes, Blueberries
Flea Beetle	Beans, Cucumbers, Melons, Squash, Peas
Aphids	

ACCEPTED
APR 17 1971
 UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 572-53 SUBJECT TO ATTACHED COMMENTS.



ROCKLAND



Net Weight One Pound

ACTIVE INGREDIENTS:

Rotenone	1%
Other Cube Extractives	2%
INERT INGREDIENTS	97%

DIRECTIONS

For best results plants should be dusted as soon as pests appear and repeated at seven day intervals. Proper application and as complete coverage as possible helps to secure successful control of insect pests. Both lower and upper sides of the plant leaves should be thoroughly dusted. The kind and age of the plants being treated will determine the amount of dust required. The quantity will vary from 1/2 pound to 1 pound per 1,000 square feet. ROCKLAND ROTENONE 1% DUST should be applied in dry weather, however, the plants must not necessarily be dry, as its effectiveness is not lost on wet plants.

The maximum kill is obtained in 1 to 3 days with effectiveness lasting a number of days, depending on weather conditions.

A good hand or power duster of the usual type is satisfactory for use.

CAUTION: Do not use on food crops within one day of harvest.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

U.S.D.A. Reg. No. 572-53