

ROCKLAND CHEMICAL CO., INC.

7/24/70
125770
CMAA 12/1/70 01205

Cameraman's Note
Poor Copy

February 10, 1970

PROPOSED LABEL COPY
FRONT PANEL

RE: C. H. L. 125770

ROCKLAND
MEMPHIS, TENNESSEE 381

1. Controls insects on ornamentals, shade and
Fruit Trees.

2. ~~substitutes for DDT~~ ~~Disinfectant~~ ~~over Broad Spectrum Insecticide~~
3. Controls adult mosquitoes, gypsy moths & others listed

~~KEEP OUT OF REACH OF CHILDREN~~
See Back Panel for complete caution statement

Manufactured by
ROCKLAND CHEMICAL CO., INC.
Passaic Avenue, N. Caldwell, N. J.

BACK PANEL

DIRECTIONS FOR USE

This is an emulsifiable concentrate
Please read entire label before using

For leaf feeding insect control on shade trees and ornamentals. Mix 4 to 6
tablespoons with 3 gallons of water (2 pints per 50 gallons water). Spray
when small larvae begin feeding. For control of cankworms, Gypsy moth,
larchworms, Juniper webworms, leaf tier, and willow, pine sawflies, spruce
sawflies on deciduous trees and conifers, direct spray to cover all foliage.
Do not over spray. Do not spray Chinese Elm, Japanese Maples, Red Maples
and Redbuds.

For Control of Fruit Tree Insects, apply only to apples, pears, peaches,
apricots, cherries and plums. Mix 4 to 6 tablespoons with 3 gallons of
water (2 pints per 50 gallons water). Spray to cover all surfaces of foliage
and fruit. Begin at petal fall and repeat at 7 to 14 day intervals as need-
ed to control apple maggot, cherry fruit worm, codling moth, leaf hoppers,
curculio, tent caterpillar and Japanese beetle. Do not apply later
than 7 days before harvest on apples, cherries, pears, plums, or later than
21 days of harvest on peaches and apricots.

Do not use with materials containing sulphur or other substances incompatible
with oil. Do not use on crops within 14 days of application of any material
containing sulphur.

For Adult Mosquito Control, Mix 1 pint with 2 gallons of water. Spray at the
rate of 1 oz. per 1000 square feet (2 gallons per acre). Apply a fine mist
spray to penetrate the areas occupied by adult mosquitoes. Apply in early
morning or at night when wind is not in excess of 3 miles per hour. Timing
best control is with the greatest mosquito activity. Repeat as necessary.

- 1. ~~Controls~~ **Fruit Trees.**
- 2. ~~controls~~ **controls adult mosquitoes, gypsy moths & others listed.**

KEEP OUT OF REACH OF CHILDREN
See Back Panel for complete caution statement

Manufactured by
ROCKLAND CHEMICAL CO., INC.
Passaic Avenue, N. Caldwell, N. J.

FRONT PANEL

DIRECTIONS FOR USE

This is an emulsifiable concentrate
Please read entire label before using

For leaf feeding insect control on shade trees and ornamentals. Mix 4 to 6 tablespoons with 3 gallons of water (2 pints per 50 gallons water). spray when small larvae begin feeding. For control of cankworms, Gypsy moth, larchworms, Juniper webworms, leaf tier and rollers, pine sawflies, spruce sawflies on deciduous trees and conifers, direct spray to cover all foliage. Do not over spray. Do not spray Chinese Elm, Japanese Maples, Red Maples and Redbuds.

For Control of fruit tree insects, apply only to apples, pears, peaches, apricots, cherries and plums. Mix 4 to 6 tablespoons with 3 gallons of water (2 pints per 50 gallons water). Spray to cover all surfaces of foliage and fruit. Begin at petal fall and repeat at 7 to 14 day intervals as needed to control apple maggot, cherry fruit worm, codling moth, leaf hoppers, rose curculio, tent caterpillar and Japanese beetle. Do not apply later than 7 days before harvest on apples, cherries, pears, plums, or later than 21 days of harvest on peaches and apricots.

Do not use with materials containing sulphur or other substances incompatible with oil. Do not use on crops within 14 days of application of any material containing sulphur.

For Adult Mosquito control, mix 1 pint with 2 gallons of water. Spray at the rate of 1 oz. per 1000 square feet (2 gallons per acre). Apply a fine mist spray to penetrate the areas occupied by adult mosquitoes. Apply in early morning or at night when wind is not in excess of 3 miles per hour. Timing should coincide with the greatest mosquito activity. Repeat as necessary. Avoid spray drift over streams, lakes and ponds. Do not apply this treatment where cooling towers are grown.

Age 1

572-5A

SEARCHED
SERIALIZED
INDEXED
FILED
FBI - NEW YORK
MAY 19 1954

ROCKLAND CHEMICAL CO., INC.

200 West 204
New York, N.Y. 10011

Camerman's Note

Poor Copy

ROCKLAND CHEMICAL CO. (NY 10011)
DATA TABLE (NY 10011)

ROCKLAND CHEMICAL CO. (NY 10011)

CAUTION: Keep out of reach of children. Do not use on humans. Do not get on skin, may cause irritation. In case of contact, wash with soap and water. Do not breathe spray mist. Do not contaminate feed or foodstuffs. Do not store or use near fire or open flame.

This product is toxic to fish. Keep out of streams, lakes or ponds. Do not contaminate water by cleaning of equipment or disposal of waste. Do not re-use empty drum. Return drum to reconditioner or destroy by perforating or crushing and burying in a safe place. This product is toxic to bees and should not be applied when bees are actively visiting the area.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NET WEIGHTS:

Methoxychlor, Technical*..... 25.00

Xylene..... 70.00

NET WEIGHTS:..... 95.00

100.00

*25.0% Methoxychlor, technical is equivalent to ~~25.0%~~ 22.0%

2,2-bis (p-Methoxyphenyl)-1,1,1-trichloroethane and

3.07% other isomers and reaction products.

contains 2 lbs. Technical Methoxychlor per gallon

REG. NO. 571-56