

Am 18 572-58 19/72

ROCKLAND
VEGETABLE
TOMATO
DUST

Use as a dust or spray.

Controls listed insects and diseases affecting
listed vegetables

1/2 to 1 pound per 1000 sq. ft. coverage.

ACTIVE INGREDIENTS:

Rotenone.....1.0%
Other Associated Resins.....2.0%
Copper as metallic (derived from
basic copper sulfate).....7.0%

INERT INGREDIENTS.....90.0%

EPA Reg. No. 572-58

EPA Est. 572-NJ-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a Physician or Poison Control Center. IF IN EYES: Flush with plenty of water. Get Medical attention. IF ON SKIN: Remove contaminated clothing and shoes. Wash affected area with soap and water. Launder clothing before reuse. IF INHALED: Remove from contaminated atmosphere.

See back panel for additional precautions.

NET WEIGHT One Pound

ACCEPTED
JUL 15 1992
572-58

ACCEPTED
JUL 15 1992
572-58

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or swallowed. Avoid breathing dust or contact with eyes or skin. Wear regular long-sleeved work clothing when applying. Wash thoroughly after handling. Food utensils such as teaspoons and cups should not be used for food after use with pesticides.

ENVIRONMENTAL HAZARDS:

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For best results, plants should be dusted and sprayed as soon as pests appear and repeated at 7 day intervals. Proper application and complete coverage helps secure successful control of insect pests. Both lower and upper sides of the plant should be treated.

Crop	Insects Controlled
BROCCOLI CABBAGE CAULIFLOWER	Cabbage Loopers, Flea Beetles, Diamond Back Moth, and Downy Mildew
CUCUMBERS MELONS SQUASH	Striped and Spotted Cucumber Beetles, Scab, Leaf Soot, Anthracnose and Bacterial Wilt
GRAPES	Black Rot and Grape Flea Beetle
CELERY EGGPLANTS POTATOES	Early and Late Blight, Anthracnose, Flea Beetles and Colorado Potato Beetle
STRAWBERRIES	Leaf Soot and Spider Bug
LIMA BEANS STRING BEANS	Mexican Bean Beetles, Downy Mildew and Anthracnose
TOMATOES	Early and Late Blight, Leaf Soot, Aphids, Fruit Worm and Flea Beetle

DUSTING: ROCKLAND VEGETABLE-TOMATO DUST is ready for use and may be dusted from the sifter can or used in a dust gun. For best results, apply in early morning or late afternoon. The quantity required will vary from 1/2 lb. to 1 lb. per 1000 sq. ft.

SPRAYING: Premix sufficient water with 6 to 8 tablespoons of ROCKLAND VEGETABLE-TOMATO DUST to form a thin, smooth paste. Add paste to water in garden sprayer to make one gallon spray. Mix thoroughly. 1 lb. makes 10 gallons of spray.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store unused material, in original container only in a locked storage area out of reach of children and animals. Keep container tightly closed.

DISPOSAL: Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

← This package is sold by weight, not by volume. Some settling of contents may have occurred during shipping and handling.

Manufactured by

 Rockland Chemical Co., Inc.
West Caldwell, NJ 07090 U.S.A.

Product A05801
EPA Reg. No. 572-58 EPA Est. 572-NJ-1

© Rockland Chemical Co., Inc. 1984

BEST AVAILABLE COPY