

ACCEPTED
 Apr 1, 1963
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR FEDERAL REGISTRATION NUMBER
 572-62



ROCKLAND

Net Weight One Pound

ACTIVE INGREDIENTS:

| | |
|---------------------------|--------|
| Malathion* | 6.0% |
| Captan ** | 6.0% |
| Methoxychlor Technical*** | 10.0% |
| Sulfur**** | 25.0% |
| INERT INGREDIENTS: | 53.0% |
| | 100.0% |

*O,O-Dimethyl dithiophosphate of diethylmercaptosuccinate.
 **N-trichloromethyl thio-ε-cyclohexene-1,2-dicarboximide.
 ***Equivalent to 8.8% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane.
 ****Not less than 98.0% passing 325 mesh.

PARTIAL LIST OF INSECTS CONTROLLED:

| | |
|----------------------------|------------------|
| ON APPLES, PEARS, PEACHES: | ON CHERRIES: |
| Aphids | Aphids |
| Cankerworm | Thrips |
| Coddling Moth | ON PLUMS: |
| Red Spider Mite | Curculio |
| Leaf Hopper | Plant Bugs |
| Leaf Roller | ON STRAWBERRIES: |
| Oriental Fruit Moth | Spittlebug |
| Tent Caterpillar | Strawberry Weev |

PARTIAL LIST OF DISEASES CONTROLLED:

| | |
|---------------------|-----------------|
| ON APPLES AND PEARS | ON STRAWBERRIES |
| Scab | Botrytis Rot |
| Powdery Mildew | |



ROCKLAND FRUIT TREE SPRAY

DIRECTIONS FOR USE

Timing is important for both disease and insect control. For best results follow a regular spray schedule. Spray thoroughly to cover all fruit and leaf surfaces. A 6 ft. to 8 ft. tree requires about 2 gallons of spray; larger trees need 6 or more gallons. Mix thoroughly before using and occasionally during use. For hose-end sprayer, follow manufacturer's directions. Do not store diluted spray.

DILUTION TABLE

| | | |
|----------------------|----|---------------------|
| Mix 4 Tablespoonfuls | to | 1 gallon of water |
| Mix 1 Cupful | to | 3 gallons of water |
| Mix 1 Pound | to | 12 gallons of water |

APPLES, PEARS: Spray when new growth appears and repeat at weekly intervals up to bloom. Then spray at 10 day intervals AFTER petals fall. Do not spray within 7 days of harvest.

PEACHES: Starting in the pink stage, make about seven spray applications at 10 day intervals. Do not apply within 21 days of harvest. For Peach Tree Borer, starting about July 1st spray or paint trunk from crotches to ground at monthly intervals using 2 lb. ROCKLAND FRUIT TREE SPRAY per gallon of water. Do not apply this concentration to fruit.

CHERRIES: Apply at full bloom, petal fall, 2 weeks after petal fall and then 2 weeks before harvest.

PLUMS: Make first application just before blooms open. Make about 7 sprayings at 10 day intervals after petals fall. Do not spray within 7 days of harvest.

STRAWBERRIES: Apply when new growth starts, then at 7 to 14 day intervals. Do not apply within 14 days of harvest.



ROCKLAND



ROCKLAND
Fruit Tree
Spray

- ① INSECT AND DISEASE CONTROL ON APPLES,
PEARS, PEACHES.
- ② CONTAINS MALATHION, METHOXYCHLOR, CAPTAN.
- ③ ONE POUND MAKES 12 GALLONS OF SPRAY.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

ROCKLAND CHEMICAL CO., Inc.
Passaic Avenue, West Caldwell, N. J.

ROCKLAND FRUIT TREE SPRAY
DIRECTIONS FOR USE