## FRUIT TREE SPRAY

	hlor, Captin. CC
ACTIVE INGREDIENTS:  Malathion*	13 % 10.00% 25.00%

\*0,0-dimethyl phosphorodithioate of diethyl mercaptoxuccinate.

\*\*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide.

\*\*\*Equivalent to 8.8% 2,2-bis(p-methoxyphenyl)-1,1,1
trichloroethane and 1.2% other isomers and related compounds.

\*\*\*\*Not less than 98.0% passing 325 mesh.

EPA REG. No. 572-62

EPA Est. No 572-NJ-1

KEEP OUT OF REACH OF CHILDREN ( 12\_pt. type)

DANGER (18 pt. type)

## PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Remove contaminated clothing and separately launder coming before reuse.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes)... Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

See Back Panel for additional precautionary statements.

Net Weight One Pound.

## STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Call a physician immediately, may cause irreversible eye damage. Flush eyes with plenty of water.

IF SWALLOWED: Call physician or poison control center immediately. Drink one or two glasses of water and induce vomiting by touching finger to back of throat. do not give anything by mouth to an unconscious erson. NOTE TO PHYSICIAN: This product may ause cholinesterase inhibition. Atropine is antidotal.

IF ON SKIN: Wash with plenty of soap and water. Launder clothes before reuse.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS: Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse. Wear chemical resistant gloves when using indoors or outdoors. When using, wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

Timing is important for both disease and insect control. For best results follow a regular spray\_schedule. Spray thoroughly to cover all fruit and leaf surfaces. A 6 ft. to 8 ft. tree requires about 3 gallons of spray; larger trees need 6 or more gallons. Mix thoroughly before using and occasionally during use. For hose-end sprayer, follow manufacturer's directions. Coarse sprays are less likely to drift. Do not apply within 10 days before or after oil spray treatment. Do not store diluted spray.

### DILUTION TABLE

Mix 5 Tablespoons to 1 gallon water. Mix 1 Cup to 3 gallons of water. Mix 1 Pound to 12 gallons of water.

APPLES, PEARS: Spray when new growth appears and repeat at weekly intervals
AFTER petals fall. Do not spray within 7 days of harvest.

PEACHES: Starting on the pink stage, make about seven spray applications.

at 10 day intervals. Do not apply within 21 days of harvest. For Peach Tree Borer, starting about July 1st, spray or paint trunk from critches to ground at monthly intervals using ½ 1b. Rockland Fruit Tree Spray per gallon of water.

Do not apply this concentration to fruit.

CHERRIES: Apply at full bloom, petal fall, 2 weeks after petal fall and then 2 weeks before harvest.

PLUMS: Nake first application just before blooms open. Make about the sprayings at 10 day intervals after petals fall. Do not spray within 7 days of harvest.

STRANBERRIES: Apply when new growth starts, then at 7 to 14 day intervals. Do not apply within 14 days of harvest.

Crop	Partial List of Insects Controlled	
Apples Pears Peaches	Aphids, Cankerworms, Coddling Moth, Red Spider Mile, Leat Hopper, Leaf Roller, Oriental Fruit Moth, Tent Caterpillar	<b>7</b>
Cherries	Aphids, Thrips	
Plums	Curculio, Plant Bugs 1	·
Strawberries .	Spittlebug, Strawberry Weevil	
Crop	. Partial List of Diseases Controlled	
Apples Pears	Scab, Powdery Mildew	
Cherries Peaches Plums	Brown Rot	
Strawberries	Botrytis Rot	

# STORAGE AND DISPOSAL

STORAGE: Store material in original container only, in a locked, dry, storage area out of reach of children and animals. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse this container. Wrap in newspaper and discard in trash.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

This package is sold by weight, not by volume. Some settling of contents may have occurred during shipping and handling.

Manufactured by:
Rockland Chemical Co., Inc.
West Caldwell, NJ 07006 USA

A06201