

111 11 11 511


1

ROCKLAND

FRUIT TREE SPRAY

.....Insect and Disease Control on Apples,
..... Contains Malathion, Methoxychlor, Captan.
..... One Pound Makes 11 Gallons of Spray.

Peaches, Cherries



ACTIVE INGREDIENTS:

Malathion ^W	5.00%
Captan ^W	5.37%
Related Compounds.....	.13%
Methoxychlor, Technical ^W	10.00%
Sulfur ^W	25.00%

INERT INGREDIENTS.....	59.00%
	100.00%

^W0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate.
^WTrichloromethylthio-4-cydonexene-1,2-dicarbonylthio.
^WEquivalent to 3.8% 1,1-dis(p-methoxyphenyl)-1,1,1-trichloroethane and 1.2% other isomers and related compounds.
~~W~~Not less than 98.0% passing 325 mesh.

EPA REG. No. 572-62

EPA Est. No 572-NJ-1

KEEP OUT OF REACH OF CHILDREN (12 pt. type)

DANGER (16 pt. type)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate resources.

See Back Panel for additional precautionary statements.

Net Weight One Pound.

BEST AVAILABLE COPY

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Call a physician immediately, may cause irreversible eye damage. Flush eyes with plenty of water.

IF SWALLOWED: Call physician or poison control center immediately. Drink one or two glasses of water and induce vomiting by touching finger to back of throat. Do not give anything by mouth to an unconscious person. NOTE TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal.

IF ON SKIN: Wash with plenty of soap and water. Launder clothing before reuse.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS: Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse. Wear chemical resistant gloves when using indoors or outdoors. When using, wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

FRUIT TREE SPRAYS: Timing is important for both disease and insect control. For best results, follow a regular spray schedule. Spray thoroughly to cover all fruit and leaf surfaces. Mix thoroughly at the following rates:

- 5 Tbsp. to 1 gallon of water.
- 1 Cup to 3 gallons of water.
- 1 lb. to 12 gallons of water.

Continue to mix occasionally during use. Apply as follows:

- 1-2 pints per small tree (6-8 ft.)
- 1-2 quarts per medium tree (8-12 ft.)
- 1-3 gallons per large tree (over 12 ft.)

For hose end sprayer, follow manufacturer's directions to apply above rates. Coarse sprays are less likely to drift. Do not apply within 10 days before or after oil spray treatment. Do not store diluted spray.

BEST AVAILABLE COPY

SUGGESTED TIMING OF SPRAY APPLICATION:

APPLES, Spray when new growth appears and repeat at weekly intervals AFTER petals fall. Do not spray within 7 days of harvest.

PEACHES: Starting on the pink stage, make about seven spray applications at 10 day intervals. Do not apply within 21 days of harvest. For Peach Tree Borer, starting about July 1st, spray or paint trunk from crotches to ground at monthly intervals using 1/2 lb. Rockland Fruit Tree Spray per gallon of water. Do not apply more than 2 gallons of mixture to any one acre of land at this dilution. Do not apply this concentration to fruit.

CHERRIES: Apply at full bloom, petal fall, 2 weeks after petal fall and then 2 weeks before harvest. Do not spray within 7 days of harvest.

PLUMS: Make first application just before blooms open. Make about 7 sprayings at 10 day intervals after petals fall. Do not spray within 7 days of harvest.

STRAWBERRIES: Apply when new growth starts, then at 7-14 day intervals. Mix 1/4 lb. Fruit Tree Spray with 3 gallons of water and spray over 225 sq. ft. of plantings. Do not apply within 14 days of harvest.

PARTIAL LIST OF INSECTS CONTROLLED:

APPLES, PEACHES: Aphids, Cankerworms, Coddling Moth, Red Spider Mite, Leaf Hopper, Leaf Roller, Oriental Fruit Moth, Tent Caterpillar.

CHERRIES: Aphids, Thrips.

PLUMS: Curculio, Plant Bugs.

STRAWBERRIES: Spittlebug, Strawberry Weevil

PARTIAL LIST OF DISEASES CONTROLLED:

APPLES Scab, Powdery Mildew.

CHERRIES, PEACHES PLUMS: Brown Rot.

STRAWBERRIES: Botrytis Rot.

STORAGE & DISPOSAL

STORAGE: Store material in original container only, in a locked, dry storage area out of reach of children and animals. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse this container. Wrap in newspaper and discard in trash.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

This package is sold by weight, not by volume. Some settling of contents may have occurred during shipping and handling.

A06291

Mfg.

Rockland Chemical Co., Inc.
West Caldwell, N.J. 07080

