

CAUTION

Avoid spray drift to susceptible plants as 2,4-D will injure cotton, beans, peas, ornamentals, etc. Coarse sprays are less likely to drift. In addition, vapors from this product may injure susceptible plants in the immediate vicinity. DO NOT use the same spray equipment for applying other materials to plants as injury will result. DO NOT store near fertilizers, seeds, insecticides, or fungicides. Avoid contact with skin, eyes or clothing.

NON-WARRANTY

Seller makes no warranty of any kind, express or implied, concerning the use of this product. Buyer assumes all risks of use or handling whether in accordance with directions or not.

Net Weight 4.00 lbs.
Gross Weight 4.10 lbs.
Net Weight 4.00 lbs.

ROCKLAND

LV-2,4-D

Weed Killer

(Low Volatile)

CONTAINS 4 LBS. 2,4-D ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS

Butoxy Ethoxy Propanol Ester of
2,4-Dichlorophenoxyacetic Acid 76.0%

INERT INGREDIENTS 24.0%

TOTAL 100.0%

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 43.2%

Contains Butoxy Ethoxy Propoxy Propanol Ester of 2,4-D.

Manufactured by

MADE IN U.S.A.

ROCKLAND CHEMICAL CO INC

PASSAIC AVENUE

35

WEST CALDWELL, N. J.

ACCEPTED
MAR 20
UNDER FEDERAL
FUNGICIDE AND
FOR ECONOMIC PL
UNDER NO. 51

ROCKLAND

LV-2,4-D Weed Killer

(Low Volatile)

CONTAINS 4 LBS. 2,4-D ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS


*Butoxy Ethoxy Propanol Ester of 2,4-Dichlorophenoxyacetic Acid	70.0
--	------

INERT INGREDIENTS	24.0
-------------------	------

TOTAL	100.0%
-------	--------


Equivalent to 2,4-Dichlorophenoxyacetic Acid	43.2
---	-------

Contains Butoxy Ethoxy Propoxy Propanol Ester of 2,4-D



NET CONTENTS

GALLONS



DIRECTIONS FOR USE

This product may be applied by either low pressure, low volume equipment or by conventional high pressure sprayer. It may be added to either water or solvents in making application.

The time of application is determined largely by the stage of growth of the weed, but when applied as a selective spray the stage of growth of the crop plant must be considered.

SMALL GRAIN CROPS Oats, Barley, Wheat, Rye. Apply 1 pint per acre on susceptible weeds as wild oat and ryegrass and 1 1/2 pints per acre on perennials such as bindweed. Make application any time after grain plants are 3-4" tall except blossom stage shall be avoided.

CORN Weeds may be controlled by application to 6 days after planting or 10 days after corn emerges through soil.

POST-EMERGENT E. J. When weeds emerge, 1/2 pint per acre on perennials, weeds will require 1 1/2 pints per acre. Apply application at or near to growing point of corn if possible.

SUGAR CANE Two 1/2 pints per acre will kill morning glory and alligator weed. Delay application until cane is two feet high.

RICE The use of 1 1/2 pints per acre kills most weeds in this crop. Consult local authorities for timing applications. **FLAX** Susceptible weed plants require 1 1/2 pints per acre. Apply when flax is 4-6" high. Be especially careful in following dosage on this crop.

PASTURES Apply 1/2 pint per acre on annual weeds and 1/2 quart per acre on perennials.

FENCE ROWS, ROADSIDES, DITCHBANKS Apply 1/2 quart per acre as soon as weeds appear and 2 quarts per acre if woody plants such as poison oak, sumac or other brush are present. Avoid contamination of water used for irrigation purposes.

UNPLANTED AREAS Apply 1 pint per acre on annuals and 1 quart per acre if perennials are a problem.

LAWNS AND GOLF FAIRWAYS Apply 1/2 pints per acre depending on weed infestation and season of year. Avoid use on bent grass and fescue.

All use quantities are based on time of application. Quantity of spray solution will depend on equipment used. For example, 1/2 gallon for low volume sprayers and 1/4 gallon for high volume equipment. Sufficient spray solution should be used to cover most susceptible weeds. For more information consult your local Rockland Sales Office or the nearest Rockland Sales Office.

ROCKLAND CHEMICAL CO INC.
 PASSAIC AVENUE 35 WEST CALDWELL, N. J. 35

CAUTION

Avoid spray drift to susceptible plants as 2,4-D will injure cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) In addition, vapors from this product may injure susceptible plants in the immediate vicinity. DO NOT use the same spray equipment for applying other materials to plants as injury will result. DO NOT store near fertilizers, seeds, insecticides, or fungicides. Avoid contact with skin, eyes or clothing.

NON-WARRANTY

Seller makes no warranty of any kind, express or implied, concerning the use of this product. Buyer assumes all risks of use or handling whether in accordance with directions or not.

Licensed under United States Patent Nos.

2,390,941	2,394,916
2,396,513	2,412,510
2,472,347	2,453,983

Re-issue 23,115

ROCK

LV-2

Weed

(Low V)

CONTAINS 4 LBS. 2,4-D AC

ACTIVE INGREDIENTS:

*Butoxy Ethoxy Propanol Ester
2,4-Dichlorophenoxyacetic A

INERT INGREDIENTS

TOTAL

*Equivalent to 2,4-Dichloroph
(Contains Butoxy Ethoxy Pro

Manufc