


NET CONTENTS 8 OUNCES



ROCKLAND



LETTUCE: To control Flea Beetle, Leafhopper, Six Spotted Leafhopper, add 1 1/2 level teaspoons to 2 1/2 gallons of water to cover 1,000 sq. ft. or 1 1/2 lbs. in 100 gallons per acre. To control Corn Earworm, Armyworm, Cabbage Caterpillar, Spittlebug, Lygus and Stink Bug, add 4 level teaspoons in 2 1/2 gallons of water to cover 100 sq. ft. or 4 lbs. in 100 gallons per acre. Do not apply later than 3 days before harvest.

ORNAMENTALS: For control of Japanese Beetle, Aphis, Bagworm, Blister Beetle, Box Elder Bug, certain Leaf Miners, Elm Leaf Beetle, Flea Beetle, Leafhopper, Leaf Roller, Orange Tortrix, Plant Bug, Scale Insects, Tent Caterpillars, Exposed Thrips, apply 2 level teaspoons per gallon of water to cover 1,000 sq. ft. or 2 lbs. per 100 gallons per acre, uniformly to Andromeda, Arbor Vitae, Ash, Aster, Azalea, Bamboo, Beech, Begonia, Birch, Boxwood, Camellia, Carnation, Ceanothus, Cedar, Chinese Elm, Chrysanthemum, Cypress, Dahlia, Delphinium, Dogwood, Elm, Euonymus, Fir, Fuchsia, Gladiolus, Hemlock, Hickory, Holly, Hornbeam, Hydrangea, Fern, Maple, Marigold, Nasturtium, Oak, Orchid, Ornamental Mulberry, Petunia, Pine, Poplar, Privet, Rhododendron, Rose, Sassafras, Spirea, Verbena, Viburnum, Willow, Wisteria, Yew, Zinnia.

NOTE: SEVIN* does not control mites but is compatible with all common insecticides. SEVIN* is toxic to honeybees. Avoid use during flowering and pollination period.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of dust. Skin contact with any insecticide may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved clothing. Change to clean clothing daily. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.** Note for physicians: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

This can is sold by weight and not by volume. Occasionally a can may not appear full due to settling of contents, but it contains full labeled weight.

WARRANTY: Seller makes no warranty of any kind, express or implied, concerning the use or handling of this material, whether in accordance with directions or not.

ACCEPTED
4-10-67
572-106
UNDER THE FEDERAL INSECT
FUNGICIDE AND RODENTICIDE
ACT, DOMESTIC POISON REG
ISTRATION NO. 10000 SU
TO ATTACH AND COMMENTS.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate*) 50%
INERT INGREDIENTS: 50%

*Trade Mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

DIRECTIONS FOR USE

ROCKLAND 50% SEVIN* Wettable Powder will control the following insects on those plants listed. For adequate control, spray coverage of plants should be uniform and complete.

ROCKLAND 50% SEVIN* Wettable Powder is compatible with commonly used miticides, fungicides and insecticides. Do not use with lime or other alkaline materials.

CHINCH BUG CONTROL ON TURF: Apply 6 ounces in 30 to 40 gallons of water per 1,000 sq. ft. of lawn.

LAWN INSECT CONTROL: To control Leaf Hoppers, Sod Webworms or Lawn Moths, Ants, Earwigs, Fleas and Chinch Bugs, apply 6 ounces in 30 to 40 gallons of water per 1,000 sq. ft. of lawn.

BEANS (Snap, Lima, Soy and Field Dry): To control Mexican Bean Beetle, add 1 level teaspoon of ROCKLAND 50% SEVIN* Wettable Powder to 2 1/2 gallons of water and apply over 1,000 sq. ft. or 1 lb. in 100 gallons per acre. To control Lygus and Stink Bugs, add 4 level teaspoons to 2 1/2 gallons per 100 sq. ft. or 4 lbs. in 100 gallons per acre.

If needed, repeat at 7 to 10 day intervals. May be applied on the day of harvest. Do not feed treated vines to dairy or meat animals.

TOMATOES, POTATOES: To control Colorado Potato Beetle and Leaf Hoppers, add 1 1/2 level teaspoons to 2 1/2 gallons of water to cover 1,000 sq. ft. or 1 1/2 lbs. in 100 gallons per acre. To control Flea Beetle, add 2 level teaspoons in 2 1/2 gallons to cover 1,000 sq. ft. or 2 lbs. in 100 gallons per acre. To control Lygus Bugs, Tomato Fruitworm, Tomato Hornworm, add 4 level teaspoons in 2 1/2 gallons of water to cover 1,000 sq. ft. or 4 lbs. in 100 gallons per acre. Repeat as needed at 7 to 10 day intervals.

CORN: To control Corn Earworm, European Corn Borer, Fall Armyworm, Japanese Beetle add 4 level teaspoons in 2 1/2 gallons of water to cover 1,000 sq. ft. or 4 lbs. in 100 gallons per acre. Do not apply to corn within 7 days of harvest for forage.

U.S.D.A. Reg. No. 572-106



ROCKLAND

NET CONTENTS 8 OUNCES



LETTUCE: To control Flea Beetle, Leafhopper, Six Spotted Leafhopper, add 1 1/2 level teaspoons to 2 1/2 gallons of water to cover 1,000 sq. ft. or 1 1/2 lbs. in 100 gallons per acre. To control Corn Earworm, Armyworm, Cabbage Caterpillar, Spittlebug, Lygus and Stink Bug, add 4 level teaspoons in 2 1/2 gallons of water to cover 100 sq. ft. or 4 lbs. in 100 gallons per acre. Do not apply later than 3 days before harvest.

ORNAMENTALS: For control of Japanese Beetle, Aphis, Bagworm, Blister Beetle, Box Elder Bug, certain Leaf Miners, Elm Leaf Beetle, Flea Beetle, Leafhopper, Leaf Roller, Orange Tortrix, Plant Bug, Scale Insects, Tent Caterpillars, Exposed Thrips, apply 2 level teaspoons per gallon of water to cover 1,000 sq. ft. or 2 lbs. per 100 gallons per acre, uniformly to Andromeda, Arbor Vitae, Ash, Aster, Azalea, Bamboo, Beech, Begonia, Birch, Boxwood, Camellia, Carnation, Ceanothus, Cedar, Chinese Elm, Chrysanthemum, Cypress, Dahlia, Delphinium, Dogwood, Elm, Euonymus, Fir, Fuchsia, Gladiolus, Hemlock, Hickory, Holly, Hornbeam, Hydrangea, Fern, Maple, Marigold, Nasturtium, Oak, Orchid, Ornamental Mulberry, Petunia, Pine, Poplar, Privet, Rhododendron, Rose, Sassafras, Spirea, Verbena, Viburnum, Willow, Wisteria, Yew, Zinnia.

NOTE: SEVIN* does not control mites but is compatible with all common insecticides. SEVIN* is toxic to honeybees. Avoid use during flowering and pollination period.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of dust. Skin contact with any insecticide may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved clothing. Change to clean clothing daily. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.** Note for physicians: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

This can is sold by weight and not by volume. Occasionally a can may not appear full due to settling of contents, but it contains full labeled weight.

WARRANTY: Seller makes no warranty of any kind, express or implied, concerning the use or handling of this material, whether in accordance with directions or not.

4-10-67
572-106

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 50%

INERT INGREDIENTS:

50%

Total 100%
Mills, Merck & Co., Inc., Kenilworth, N.J.
© 1967, Merck & Co., Inc.

DIRECTIONS FOR USE

ROCKLAND 50% SEVIN* Wettable Powder will control the following insects on those plants listed. For adequate control, spray coverage of plants should be uniform and complete.

ROCKLAND 50% SEVIN* Wettable Powder is compatible with commonly used miticides, fungicides and insecticides. Do not use with lime or other alkaline materials.

CHINCH BUG CONTROL ON TURF: Apply 6 ounces in 30 to 40 gallons of water per 1,000 sq. ft. of lawn.

LAWN INSECT CONTROL: To control Leaf Hoppers, Sod Webworms or Lawn Moths, Ards, Earwigs, Fleas and Chinch Bugs, apply 6 ounces in 30 to 40 gallons of water per 1,000 sq. ft. of lawn.

BEANS (Snap, Lima, Soy and Field Dry): To control Mexican Bean Beetle, add 1 level teaspoon of ROCKLAND 50% SEVIN* Wettable Powder to 2 1/2 gallons of water and apply over 1,000 sq. ft. or 1 lb. in 100 gallons per acre. To control Lygus and Stink Bugs, add 4 level teaspoons to 2 1/2 gallons per 100 sq. ft. or 4 lbs. in 100 gallons per acre.

If needed, repeat at 7 to 10 day intervals. May be applied on the day of harvest. Do not feed treated vines to dairy or meat animals.

TOMATOES, POTATOES: To control Colorado Potato Beetle and Leaf Hoppers, add 1 1/2 level teaspoons to 2 1/2 gallons of water to cover 1,000 sq. ft. or 1 1/2 lbs. in 100 gallons per acre. To control Flea Beetle, add 2 level teaspoons in 2 1/2 gallons to cover 1,000 sq. ft. or 2 lbs. in 100 gallons per acre. To control Lygus Bugs, Tomato Fruitworm, Tomato Hornworm, add 4 level teaspoons in 2 1/2 gallons of water to cover 1,000 sq. ft. or 4 lbs. in 100 gallons per acre. Repeat as needed at 7 to 10 day intervals.

CORN: To control Corn Earworm, European Corn Borer, Fall Armyworm, Japanese Beetle add 4 level teaspoons in 2 1/2 gallons of water to cover 1,000 sq. ft. or 4 lbs. in 100 gallons per acre. Do not apply to corn within 7 days of harvest for forage.

TOMATO POTATO EGG PLANT PEPPER	Colorado Potato Beetle, Flea Beetle, Leafhoppers, Tomato Fruit Worm, Fall Armyworm, Tomato Horn Worm, European Corn Borer, Stink Bug and Lygus Bugs.	20 lbs.	8 oz.	Day of Harvest
LETTUCE	Flea Beetles, Leafhoppers.	20-40 lb.	1/2 - 1 lb.	Day of Harvest
	Six-spotted Leafhopper. Corn Earworm, Armyworm, Cabbage Caterpillar, Spittlebug, Lygus and Stink Bugs	10-20 lbs. 20-30 lbs.	4 - 8 oz. 8 - 12 oz.	3 Days Before Harvest 3 Days Before Harvest
LEAF LETTUCE	As Above	As Above	As Above	3 Days Before Harvest 14 Days Before Harvest
FLOWERS and ORNAMENTALS	Bagworm, Japanese Beetles, Flea Beetles, Lace Bugs, Leafhoppers, Scale Insects, Rose Aphid, Tent Caterpillars.	20-40 lbs.	8 oz. - 1 lb.	No Time Limit

Carbaryl injures Boston ivy and Virginia creeper. Do not use this product on or near these plants.

DOGS AND CATS Brown Dog Tick, Fleas. Rub in skin and apply in sleeping quarters weekly. Do not treat kittens under 4 weeks old.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of dust. **SKIN CONTACT MAY BE HARMFUL.** Avoid contact. Wash hands before eating. Take shower or bath after work. **WEAR REGULAR LONG SLEEVED WORK CLOTHING.** Change to clean clothing daily. Note to physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



EPA Reg No. 572-107
EPA Est. 572-NJ-1

ACTIVE INGREDIENTS:
Carbaryl (1-naphthyl N-methylcarbamate*) 5%
INERT INGREDIENTS: 95%

*Trade Mark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

DIRECTIONS FOR USE

ROCKLAND 5% SEVIN* DUST is finely ground to give a superior dusting quality. To control insects listed below, dust at the recommended rates, making certain to cover foliage uniformly and completely paying particular attention to the under-sides of the leaves.

CROP	INSECTS CONTROLLED	RATE OF APPLICATION for ROCKLAND 5% SEVIN* DUST		DATE OF LAST APPLICATION
		Per Acre	Per 1000 sq. ft.	
BEANS	Mexican Bean Beetle.	10 lbs.	4 oz.	Day of Harvest
	Leaf Hoppers, Bean Leaf Beetle, Flea Beetles, Japanese Beetles.	20 lbs.	8 oz.	Day of Harvest
	Corn Ear Worm, Armyworms.	30 lbs.	12 oz.	Day of Harvest
	Lygus and Stink Bugs.	40 lbs.	1 lb.	Day of Harvest
CORN	Corn Ear Worm, European Corn Borer, Fall Armyworm, Japanese Beetle.	40 lbs.	1 lb.	Day of Harvest*

For Corn Ear Worm, apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals as infestation warrants. Re-apply immediately after rain.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

*Except 7 days before when forage is to be fed to livestock.

ACCEPTED WITH COMMENTS