

572-107  
4-25-1971


1/2

TOMATO POTATO EGG PLANT PEPPER	Colorado Potato Beetle, Flea Beetle, Leafhoppers, Tomato Fruit Worm, Fall Armyworm, Tomato Horn Worm, European Corn Borer, Stink Bug and Lygus Bugs.	20 lbs.	8 oz.	Day of Harvest
LETTUCE	Flea Beetles, Leafhoppers, Six-spotted Leafhopper, Corn Earworm, Armyworm, Cabbage Caterpillar, Spittlebug, Lygus and Stink Bugs.	20-40 lbs.	1/2 - 1 lb.	Day of Harvest
		10-20 lbs.	4 - 8 oz.	3 Days Before Harvest
		20-30 lbs.	8 - 12 oz.	3 Days Before Harvest
LEAF LETTUCE	As above	As above	As above	14 Days Before Harvest
FLOWERS AND ORNAMENTALS	Bagworm, Japanese Beetles, Flea Beetles, Lace Bugs, Leafhoppers, Scale Insects, Rose Aphid, Tent Caterpillars.	20-40 lbs.	8 oz. - 1 lb.	No Time Limit
DOGS AND CATS	Brown Dog Tick, Fleas	Rub in skin and apply in sleeping quarters weekly. Do not treat kittens under 4 weeks old.		

**CAUTION: HARMFUL IF INHALED OR SWALLOWED.** Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of dust. SKIN CONTACT WITH ANY INSECTICIDE MAY BE HARMFUL. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. WEAR REGULAR LONG SLEEVED WORK CLOTHING. Change to clean clothing daily. Note to physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

This product is toxic to bees. Any use of the product when honey bees are visiting the crop will seriously reduce bee populations. Bees are important pollinators and are essential to the production of many crops. Before using this product during such periods consult your State Agricultural Extension Service.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



U.S.D.A. Reg. No. 572-107

ACTIVE INGREDIENTS:  
Carbaryl (1-naphthyl N-methylcarbamate\*) . . . . . 5%  
INERT INGREDIENTS: . . . . . 95%

**DIRECTIONS FOR USE**

ROCKLAND 5% SEVIN\* DUST is finely ground to give a superior dusting quality. To control insects listed below, dust at the recommended rates, making certain to cover foliage uniformly and completely paying particular attention to the undersides of the leaves.

CROP	INSECTS CONTROLLED	RATE OF APPLICATION for ROCKLAND 5% SEVIN* DUST		DATE OF LAST APPLICATION
		Per Acre	Per 1000 sq. ft.	
BEANS	Mexican Bean Beetle, Leaf Hoppers, Bean Leaf Beetle, Flea Beetles, Japanese Beetles.	10 lbs.	4 oz.	Day of Harvest
	Corn Ear Worm, Armyworms.	20 lbs.	8 oz.	Day of Harvest
	Lygus and Stink Bugs.	30 lbs.	12 oz.	Day of Harvest
		40 lbs.	1 lb.	Day of Harvest
CORN	Corn Ear Worm, European Corn Borer, Fall Armyworm, Japanese Beetle.	40 lbs.	1 lb.	Day of Harvest*

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

\*Except 7 days before when forage is to be fed to livestock.

SECRET

ACCEPTED  
APR 25 1971  
UNDER THE ELECTRONIC INFORMATION  
FUNGICIDE ACT  
FOR ELECTRONIC INFORMATION REGISTERED  
ED UNDER REG 572-107 SUBJECT  
TO ATTACHED COMMENTS.