




ROCKLAND

Net Contents 1 pint



1
2
3

ACCEPTED
4-28-67
572-123

Net Contents: 1 Pint

DIRECTIONS

ROCKLAND LAWN WEED KILLER will give the most effective control when applied to vigorously growing weeds in Spring or Fall after a rain. Apply on a warm day when there is no wind, preferably in the morning or evening. Measure the area to be sprayed and mix the correct dosage, using a conventional lawn sprayer or a sprayer with hose attachments. For requirements different than listed below, consult local turf authority.

LAWN WEEDS: For Dandelion, Plantain, Chickweed, Knotweed, Purslane, Clover, Pennywort and many others, mix 4 tablespoonfuls (2 fl. oz.) in 3 gallons of water and apply to 600 sq. ft. of lawn. Do not use on Dichondra, Lippia or Clover lawns. Do not use on new lawns until after at least second cutting. Do not water for 24 hours or mow for at least 3 weeks. Wild onion may require repeated application in late fall and early spring. For St. Augustine or Bent Grass Lawns, make 3 applications at 10 day intervals, using 1/2 strength, or 1 fl. oz., in 3 gallons of water for 600 sq. ft.

HARD-TO-KILL WEEDS: For Ragweed, Dock, Morning Glory on Lawns, Thistle along Fence Rows, Right-of-Ways, spray in the spring when leaves are full grown, using 3 tablespoons or 1 1/2 fl. oz. per gallon of water (1:100 dilution). Spray to wet all plant surfaces thoroughly. Repeat as necessary.

For control of **POISON IVY** and **WOODY PLANTS** in Wastelands, Fence Rows, **HONEYSUCKLE**, **SUMAC**, **BRAMBLES** and Noxious Weeds, use 4 fl. oz. (1/2 cup) per gallon of water. Spray to wet all plant surfaces thoroughly. Application may be repeated in 4 weeks.

CAUTION: Harmful if swallowed. Avoid inhalation and contact with skin, eyes or clothing. Wash hands after using. Keep containers closed. Do not store near heat or open flame, nor seeds, fertilizers, insecticides or fungicides. Do not contaminate water used for domestic or irrigation purposes. Destroy container when empty. Do not reuse. Keep out of reach of children. Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, grapes, ornamentals, etc. (coarse sprays are less likely to drift). Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (or do not use same spray equipment for other purposes). Vapors from the product may injure susceptible plants in the immediate vicinity.

WARRANTY: Seller makes no warranty of any kind, express or implied, concerning the use of this material. Buyer assumes all risk of use or handling, whether in accordance with directions or not.

ACTIVE INGREDIENTS:

- * Ethylhexyl ester of 2,4-dichlorophenoxyacetic Acid 15.2%
- ** Butoxyethanol Ester of Silvex (2-(2,4,5-trichlorophenoxy) Propionic acid) 7.1%

INERT INGREDIENTS:

- * Equivalent to 10% 2,4-dichlorophenoxyacetic acid or .8 lb. per gallon.
- ** Equivalent to 5% Silvex or .4 lb. per gallon.

USDA Reg. No. 572-123


ROCKLAND

NET WT. 12 LBS.

CRABGRASS PREVENTER




FOR PRE-EMERGENCE CONTROL OF CRABGRASS AND OTHER ANNUAL WEEDS AND GRASSES IN ESTABLISHED ORNAMENTAL PLANTINGS.





Net Contents: 1 pint

ROCKLAND



ROCKLAND

Lawn Weed
Killer

1
2
3

4-28-67
572-123

Net Contents: 1 Pint

DIRECTIONS

ROCKLAND LAWN WEED KILLER will give the most effective control when applied to vigorously growing weeds in Spring or Fall after a rain. Apply on a warm day when there is no wind, preferably in the morning or evening. Measure the area to be sprayed and mix the correct dosage, using a conventional lawn sprayer or a sprayer with hose attachments. For requirements different than listed below, consult local authority.

LAWN WEEDS: For Dandelion, Plantain, Chickweed, Knotweed, Purslane, Spurge, Pennycress and many others, mix 4 tablespoons (2 fl. oz.) in 3 gallons of water and apply to 600 sq. ft. of lawn. Do not use on Dichondra, Ligia or Clover lawns. Do not use on new lawns until after at least second cutting. Do not water for 24 hours or reseed for at least 3 weeks. Wild onion may require repeated application in late fall and early spring. For St. Augustine or Bent Grass Lawns, make 3 applications at 10 day intervals using 1/2 strength or 1 fl. oz. in 3 gallons of water for 600 sq. ft.

HARD-TO-KILL WEEDS: For Ragweed, Dock, Morning Glory, in Lawns, Thistle along Fence Rows, Right of Ways, Spray in the spring when leaves are full grown using 3 tablespoons or 1 1/2 fl. oz. per gallon of water (1:100 dilution). Spray to wet all plant surfaces thoroughly. Repeat as necessary.

For control of POISON IVY and WOODY PLANTS in Waterlans, Fence Rows, HONEYBUCKLE, SUMAC, BRAMBLES and Noxious Weeds, use 4 fl. oz. (1/2 cup) per gallon of water. Spray to wet all plant surfaces thoroughly. Application may be repeated in 4 weeks.

CUTION: Harmful if swallowed. Avoid inhalation and contact with skin, eyes or clothing. Wash hands after using. Keep containers closed. Do not store near heat or open flame, nor seeds, fertilizers, insecticides or fungicides. Do not contaminate water used for domestic or irrigation purposes. Destroy container when empty. Do not reuse. Keep out of reach of children. Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, grapes, ornamentals, etc. (coarse sprays are less likely to drift). Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (do not use same spray equipment for other purposes). Vapor from the product may injure susceptible plants in the immediate vicinity.

WARRANTY: Seller makes no warranty of any kind, express or implied, concerning the use of this material. Buyer assumes all risk of use or handling, whether in accordance with directions or not.

ACTIVE INGREDIENTS:

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 15.2%
**Butoxyethanol Ester of Silvex 2-(2,4,5-trichlorophenoxy) Propionic acid 7.1%

INERT INGREDIENTS

*Equivalent to 2,4-Dichlorophenoxyacetic acid or 2,4,5-trichlorophenoxy

**Equivalent to 5% Silvex or 4 lb. per gallon

USDA Reg. No. 572-123