

SEE BACK PANEL FOR COMPLETE CAUTION STATEMENT. ACTIVE INGREDIENTS:

*Ronnel (0,0-Dimethyl 0-(2,4,5-trichlorophenyl) Aromatic Petroleum derivative solvent	phosphorothioate)	24.0% 68.0%
INERT INGREDIENTS:	. .	8.0%

*From technical grade; not suitable for internal use in animals. U. S. Pat. No. 2599516.

Manufactured for

I B A, INC. PESTICIDE DIVISION Millbury Massachusetts I.B.A. RESIDUAL FLY SPRAY is to be mixed with water to make a 1% solution to be applied to Farm Buildings including Poultry Houses, Livestock Shelters, Milk Rooms, Dairy and Hog Barns for 4 to 6 week Fly Control.

As a Livestock Insect Spray, dilute with water only. Spray entire animal to wet coat thoroughly. Repeat its necessary but not oftener than every 2 weeks.

water.

100.00%

CATTLE, CAL' éS: To control Hornflies, use | quart with 121/2 gallons of water. To control Lice, use I quart with 25 gallons of water. To control Lone Star Ticks and Winter Ticks, use I quart with 8 gallons of water.

SHEEP, GOATS, LAMBS and KIDS: To control Wool Maggots, Hornflies, use I quart with 121/2 galons of water. To control Lice and Sheep Keds, use I quart with 25 gallons of water

For Cattle Backrubbers — for Hornflies and to aid in control of Face Flies if backrubbers are placed so animal can rub face. Mix 1 quart 1.8.A. RESIDUAL FLY SPRAY with 7 gallons of diesel, or similar, oil. Saturate fibre; recharge as necessary. Do not mix with used motor oil.

DIRECTIONS FOR USE

To make a 1% solution mix 1 quart of I.B.A. RESIDUAL FLY SPRAY with 6 gallons of water. Add 3 ounces of sugar for added effectiveness.

Use 1 gallon of diluted spray per 500-1000 sq. ft. Spray Walls, Ceilings, Partitions to obtain complete coverage with no run off. To prevent contamination, keep milk and milking equipment covered while spraying.

For Caged Layers on Wire, mix 1 quart with 5 gallons of water; spray poultry droppings beneath cages.

HOGS: To control Lice, use 1 quart with 25 gallons of

WARNING: Do not use on animals receiving an organic phosphate treatment from any other source at the time of treatment.

Do not use on animals receiving an organic phosphate treatment from any other source at the time of treatment.

NOTE TO PHYSICIAN: This marerial is a cholinesterase inhibitor. Treat symptomatically. Acopine is an antidate.

CAUTION: INJURIOUS TO EYES. IRRITATING TO THE SKIN AND EYES. HARMFUL IS SWALLOWED. Do not get in the eyes. Avoid contact with kin. Avoid breathing of spray mist. Wash thoroughly with scap and water after handling. Keep material out of reach of children and animals. Avoid storage near or contamination of food, feedstuffs and water supplies. Keep container closed.

COMBUSTIBLE LIQUID: Keep away from heat and open flame.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waster.

Do not reuse empty container. Destrc, it by perforating or crushing. Bury or discard in a safe place away from water supplies.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

> EPA Reg. No. 572-132 I.B.A. Stock No. 4919