

ACTIVE INGREDIENTS

CAUTION

KEEP OUT OF REACH OF CHILDREN  
DO NOT APPLY TO FACE, EYES, OR MOUTH



**ROCKLAND CHEMICAL CO., INC.**

WEST CALDWELL, N. J.

TREATS 2,500 30. FT.  
(50 x 50)



**ROCKLAND**

**CRABGRASS**

**PREVENTER**



50 ft.). Use any lawn spreader in good condition. Do not spread over larger area than recommended as such application would give poor results. To control annual bluegrass or other erratic or late-germinating grasses, a late summer or fall application of 12 lbs. per 1700 sq. ft. should supplement the early spring application (contents of bag will treat 1600 sq. ft.).

**HOW TO APPLY:** Before making application, lawn should be raked to remove extraneous material and insure uniform application. In order to obtain expected results, ROCKLAND CRABGRASS PREVENTER should be evenly applied with no skips or misses. It is suggested that the area be covered twice; the second time at right angles to the first to insure thorough coverage (See following table).

SPREADER	WIDTH OF SPREADER	APPROXIMATE SETTINGS	
		Double (twice over) Application	SINGLE (once over) Application
Lawn Beauty	18"	6 3/4	8 1/2
Scott	18"	4 1/2	8 1/2
Jackson	20"	8 1/2	11
Applegate	20"	8 1/2	11
Buch	18"	1 7/8	2 5/8
Agrico Brush On	20"	7 1/2	10 1/2

Do not rake the lawn or disturb soil surface after application.

Water in lightly if no rain falls within a week after application.

**RE-SEEDING:** When some spring seeding is necessary, there should be a delay of approximately 60 days after the application of ROCKLAND CRABGRASS PREVENTER. This delayed seeding improves the survival of desirable grass seedlings. If re-seeding is done before 60 days after application, the amount of seed should be increased 50% to insure thrifty stands. An early spring application may be made to newly seeded lawns after they have exhibited greening of newly-sprouted lawn grasses.

**WEED CONTROL IN ORNAMENTALS:** ROCKLAND CRABGRASS PREVENTER can be applied to a wide range of nursery stock for the control of the same annual weeds as listed under general information. Apply at the same rate as for turf. ROCKLAND CRABGRASS PREVENTER is specifically recommended for weed control in most ornamentals. However, do NOT use on the following plants: Bugleweed, Button pink, Carnation, Geum, Germander, Mesembryanthum, Pansy, Phlox, Sweet William, Telanthera.

**CAUTION:** Keep out of reach of children. Do not use on dichondra lawns or on newly seeded lawns. Do not store near seeds, fertilizers and other pesticides. Do not contaminate feed or foodstuffs. Do not take internally. Avoid contact with eyes. Wash after using. Fill spreader on non-turf area to avoid spilling chemical on lawn. Store in dry place.

USDA Reg. No. 572-147

**Polywall**

FREIGHT SHIPPING BAG  
MEETING REQUIREMENTS OF  
APPLICABLE FREIGHT CLASSIFICATION  
GUARANTEED BY

# Rockland

CHEMICAL CO., INC.

Products for the Lawn, Garden, Farm, Home

PASSAIC AVENUE, WEST CALDWELL, N. J.

mailing address  
P.O. BOX 207 CALDWELL, N. J. 07006  
telephone (201) 226-5151

December 3, 1970


PROPOSED LABEL COPY

FRONT PANEL:

ROCKLAND  
CRABGRASS AND LAWN WEED KILLER

Treats 5000 Square Feet

1. Kills Crabgrass, Dandelion, Plantain, Chickweed, Pigweed, and similar lawn weeds.
2. Apply late Spring, Summer or early Fall.
3. Contains DSMA and 2,4-D



CAUTION  
KEEP OUT OF REACH OF CHILDREN

See Back Panel for Complete Caution  
Read Entire Label Before Using This Product

ACTIVE INGREDIENTS:

Disodium methanearsonate..... 4.5%  
2,4-dichlorophenoxyacetic acid, Sodium Salt..... .9%  
(2,4-dichlorophenoxyacetic acid equivalent 0.855%)

INERT INGREDIENTS:.....94.6%

Net Wt. 10 lbs.

Manufactured by  
ROCKLAND CHEMICAL CO., INC.  
West Caldwell, N. J. 07006

BACK PANEL:

ROCKLAND  
CRABGRASS AND LAWN WEED KILLER  
DSMA plus 2,4-D

Water area to be treated thoroughly with hose immediately before application so that the material will adhere to the leaves of the weeds.

For best results on Crabgrass, make 2 or 3 applications 5 to 7 days apart.

For broad-leaved weeds, 1 or 2 applications may be sufficient.

Can be applied anytime weeds and Crabgrass are growing -- late Spring, Summer or early Fall.

Should new weed or Crabgrass infestation appear, repeat treatment.

Do not use on newly seeded lawns until mowed twice.

Do not apply during extreme drought as temporary browning on fescues and bentgrasses may occur. This product may injure Carpetgrass, St. Augustine grass, Dichondra and Clover in lawns.

For best results, do not reseed lawn until 2 weeks after application. Normal lawn sprinkling may be resumed 48 hours after treatment. ROCKLAND DSMA plus 2,4-D will be absorbed by weeds within 24 hours.

# Rockland

CHEMICAL CO., INC.

Products for the Lawn, Garden, Farm, Home

mailing address:  
P. O. BOX 201 CALDWELL, N. J. 07006  
Telephone: (201) 225-5151

December 3, 1970

PROPOSED LABEL COPY (CONT)  
BACK PANEL (CONT)

ROCKLAND CRABGRASS AND LAWN WEED KILLER

RECOMMENDED SPREADER SETTINGS

Scott---5

Lawn Beauty---6

Jackson---6½

Patco---6½

NOTE: Above spreader settings are approximate. Allowances must be made for condition and proper adjustment of spreader.

CAUTION: Harmful if swallowed. Avoid breathing of the dust and contact with skin, eyes or clothing. Keep away from children, domestic animals, feed or food products. Do not contaminate irrigation ditches or water used for domestic purposes. Wash hands thoroughly after using.

ANTIDOTE: If taken internally, induce vomiting and call a physician at once.

Keep livestock off treated areas.

Do not store near fertilizer, seeds, insecticides or fungicides. Avoid getting material on garden shrubs or flowers that may be injured by 2,4-D. Apply on a calm day.

WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

U.S.D.A. Reg. No. 572-148

**ERc**  
**OCKLAND**

**495**



**WEED**

**2**

**FEEED**

**5000 SQUARE FEET**

# Rockland

CHEMICAL CO., INC.



December 3, 1970

PROPOSED LABEL COPY  
FRONT PANEL:

ROCKLAND  
CRABGRASS AND LAWN WEED KILLER  
Treats 5000 Square Feet

1. Kills Crabgrass, Dandelion, Plantain, Chickweed, Pigweed, and similar lawn weeds.
2. Apply late Spring, Summer or early Fall.
3. Contains DSMA and 2,4-D

ACCEPTED  
DEC 3 1970  
10/20/70

CAUTION  
KEEP OUT OF REACH OF CHILDREN  
See Back Panel for Complete Caution  
Read entire Label Before Using This Product

ACTIVE INGREDIENTS:  
Disodium methanearsonate..... 4.5%  
2,4-dichlorophenoxyacetic acid, Sodium Salt..... .9%  
(2,4-dichlorophenoxyacetic acid equivalent 0.855%)  
INERT INGREDIENTS:.....94.6%  
Net wt. 10 lbs.

Manufactured by  
ROCKLAND CHEMICAL CO., INC.  
West Caldwell, N. J. 07000

BACK PANEL:

ROCKLAND  
CRABGRASS AND LAWN WEED KILLER  
DSMA plus 2,4-D

Water area to be treated thoroughly with hose immediately before application so that the material will adhere to the leaves of the weeds.  
For best results on Crabgrass, make 2 or 3 applications 5 to 7 days apart.  
For broad-leaved weeds, 1 or 2 applications may be sufficient.  
Can be applied anytime weeds and Crabgrass are growing -- late Spring, Summer or early Fall.  
Should new weed or Crabgrass infestation appear, repeat treatment.  
Do not use on newly seeded lawns until mowed twice.  
Do not apply during extreme drought as temporary browning on fescues and bentgrasses may occur. This product may injure Carpetgrass, St. Augustine grass, Dichondra and Clover in lawns.  
For best results, do not reseed lawn until 2 weeks after application. Normal lawn sprinkling may be resumed 48 hours after treatment. ROCKLAND DSMA plus 2,4-D will be absorbed by weeds within 24 hours.

# Rockland

CHEMICAL CO., INC.



December 3, 1970

PROPOSED LABEL COPY (CONT)  
BACK PANEL (CONT)

ROCKLAND CRABGRASS AND LAWN WEED KILLER

### RECOMMENDED SPREADER SETTINGS

Scott---

Lawn Beauty---6

Jackson---6½

Patco---6½

NOTE: Above spreader settings are approximate. Allowances must be made for condition and proper adjustment of spreader.

CAUTION: Harmful if swallowed. Avoid breathing of the dust and contact with skin, eyes or clothing. Keep away from children, domestic animals, feed or food products. Do not contaminate irrigation ditches or water used for domestic purposes. Wash hands thoroughly after using.

ANTIDOTE: If taken internally, induce vomiting and call a physician at once.

Keep livestock off treated areas.

Do not store near fertilizer, seeds, insecticides or fungicides. Avoid getting material on garden shrubs or flowers that may be injured by 2,4-D. Apply on a calm day.

### WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

U.S.D.A. Reg. no. 572-148