


624

FREIGHT SHIPPING BAG
MEETING REQUIREMENTS OF
APPLICABLE FREIGHT CLASSIFICATION
GUARANTEED BY
BEMIS COMPANY, INC. EAST PEPPERELL, MASS.

Polywall

TREATS **5000** SQ. F



ROCKLAN

WEED

&

FEED

Fertilizes your lawn and controls weeds such as Dandelion, Buckhorn, Plantain, Burdock, Pigweed, Cocklebur, Ragweed, Chickweed and many other lawn weeds.

Use anytime during growing season when weeds are present. Covers 5,000 sq. ft. at the following spreader settings:

SUGGESTED SPREADER SETTINGS

PATCO	6	JACKSON	5½
LAWN BEAUTY	5½	SCOTT	4½
MODERN	5½	CYCLONE	3

Fertilizes your lawn and controls weeds such as Dandelion, Buckhorn, Plantain, Burdock, Pigweed, Cocklebur, Ragweed, Chickweed and many other lawn weeds.

Use anytime during growing season when weeds are present. Covers 5,000 sq. ft. at the following spreader settings:

SUGGESTED SPREADER SETTINGS

PATCO	6	JACKSON.....	5½
LAWN BEAUTY.....	5½	SCOTT.....	4½
MODERN.....	5½	CYCLONE.....	3

For smaller areas, use proportionate amounts – approximately ¾ lbs. per 1,000 square feet.

DIRECTIONS

Mow lawn before applying. Apply as it comes from the package. Do not mix with other materials. Do not water after application. A light dusting, broadcast by hand or spreader is all that is necessary. If a spreader is used, wash it thoroughly before re-use.

Apply on a calm day, to avoid drifting of material as vapors from this product may injure susceptible plants in the immediate vicinity. Do not apply too close to flowers, shrubs, evergreens, etc. Be careful no application is made directly on them or over their root spread. Clover or creeping bentgrass present in lawns may be injured.

NOTE: Best results will be obtained if material is applied when grass is moist, as from morning or evening dew. Rain or watering within 24 hours after application will reduce effectiveness. Do not mow lawn until three days after treatment. Not recommended for use on newly seeded areas.

Some weeds are more resistant than others. If a repeat application is necessary, it can be made two weeks after the first application. However, lawn must be thoroughly watered before this second treatment. Areas may be re-seeded and fertilized 10 to 14 days after last application.


CAUTION: Avoid inhalation, contamination of foodstuffs and contact with skin, eyes or clothing. Wash hands after handling. Do not store near, nor allow loose material to contact seeds, fungicides, insecticides, other fertilizers. Avoid contamination of irrigation ditches or water used for domestic purposes.

624

FREIGHT SHIPPING BAG
REQUIREMENTS OF
APPLICABLE FREIGHT CLASSIFICATION
SEMI-COMMERCIAL - EAST OF THE RIVER, MASS.

Polywali

TREATS **5000** SQ. F



ROCKLAN

WEED



FEED

Fertilizes your lawn and controls weeds such as Dandelion, Buckhorn, Dandelion, Black Medweed, Cocklebur, Ragweed, Chickweed and many other weeds.

5.000

	INGRS
AWN BEA	5.00
MIXED	4.00
ACKSON	3.00
SCOTT	2.00
ONE	1.00

Apply 5 lbs. per 5000 sq. ft. of lawn.

DIRECTIONS

Apply to lawn in early April or late March. Do not mix with other fertilizers.

FEED

Fertilizes your lawn and controls weeds such as Dandelion, Buckhorn, Plantain, Burdock, Pigweed, Cocklebur, Ragweed, Chickweed and many other lawn weeds.

Use anytime during growing season when weeds are present. Covers 5,000 sq. ft. at the following spreader settings:

SUGGESTED SPREADER SETTINGS

PATCO	6	JACKSON	5½
LAWN BEAUTY	5½	SCOTT	4½
MODERN	5½	CYCLONE	3

For smaller areas, use proportionate amounts - approximately 3¼ lbs. per 1,000 square feet.

DIRECTIONS

Mow lawn before applying. Apply as it comes from the package. Do not mix with other materials. Do not water after application. A light dusting, broadcast by hand or spreader is all that is necessary. If a spreader is used, wash it thoroughly before re-use.

Apply on a calm day, to avoid drifting of material as vapors from this product may injure susceptible plants in the immediate vicinity. Do not apply too close to flowers, shrubs, evergreens, etc. Be careful no application is made directly on them or over their root spread. Clover or creeping bentgrass present in lawns may be injured.

NOTE: Best results will be obtained if material is applied when grass is moist, as from morning or evening dew. Rain or watering within 24 hours after application will reduce effectiveness. Do not mow lawn until three days after treatment. Not recommended for use on newly seeded areas.

Some weeds are more resistant than others. If repeat application is necessary, apply to the same area 10 to 14 days after the first application. Lawns may be re-seeded and fertilized 10 to 14 days after last application.

CAUTION: Avoid contamination, contamination of foodstuffs and contact with skin, eyes or clothing. Wash hands after handling. Do not store near, nor allow loose material to contact seeds, fungicides, insecticides, other fertilizers. Avoid contamination of irrigation ditches or water used for domestic purposes.

ER

\$495

OCKLAND

WEED

&

FEEED

TREATS

5000 SQUARE FEET



TREATS
5000 SQUARE FEET

- ① Weeds with 2,4-D and Silvex to control most lawn weeds.
- ② Feeds with 18-6-6 Fertilizer.
- ③ For Best results mow lawn before applying.

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR COMPLETE CAUTION
READ ENTIRE LABEL BEFORE USING THIS PRODUCT

ACTIVE INGREDIENTS

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid 40%
Isooctyl ester of Silvex 2-(2,4,5-Trichlorophenoxy)propionic acid 50%

INERT INGREDIENTS:

97.90%
100.00%

GUARANTEED ANALYSIS

Total Nitrogen 18%
Available Phosphorus 6%
Water Soluble Potash 6%
Potential Acidity equivalent to 675 lbs. Calcium Carbonate per ton.

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 1.17%
**Equivalent to Silvex 0.50%

MANUFACTURED BY

NET WT. 16 LBS

ROCKLAND CHEMICAL CO., INC.

WEST CALDWELL, N. J.