

**ACTIVE INGREDIENT:**

Technical Chlordane\*

10.0%

**INERT INGREDIENTS:**

90.0%

\*Equivalent to 6.0% Octachloro-4,7-Methanotetrahydroindane and 4.0% related compounds, on a weight basis.

**"HOMEOWNER" 10% CHLORDANE DUST**

**DIRECTIONS FOR USE**

To Make Spray Solution: Mix 2 tablespoons with a little water to make a thin paste. Stir paste into 1 gallon of water. Agitate while using. For Hose-End Sprayers follow manufacturer's directions.

To control Ants, Leaf Miners, Leaf Rollers, Thrips, Plant Bugs, Leaf Beetles, Lace Bugs, Japanese Beetles (Adults) and certain Caterpillars on FLOWERS AND ORNAMENTALS: Dust or spray all plant surfaces in the early morning or evening when the air is still. For Taxus Weevils, treat lower branches and soil around the base of the plants. For Earwigs, Crickets and Mole Crickets, use 1 pound for 700 sq. ft. of soil around plants. Repeat as necessary.

For Japanese Beetle Grubs, White Grubs and Chafer Beetle Grubs in LAWNS, apply 1 pound to 700 sq. ft. of lawn area and water thoroughly to carry insecticide into soil. For Lawn Moths, Sod Webworms, Mole Crickets and Crickets, water lightly after application. Keep children and pets off treated area until thoroughly dry. May be harmful to birds feeding on treated areas.

USE AS DUST OR SPRAY  
FOR LAWNS AND GARDENS



**ROCKLAND**

Net Weight One Pound



In VEGETABLE GARDEN, for Flea Beetles and Hornworms on tomatoes, Flea Beetles on beans, black-eyed peas, eggplant and cabbage, Cucumber Beetle, Squash Bug and Squash Vine Borer on cucumbers, melons and squash, dust 4 to 6 oz.


CHLORDANE 10% DUST per 1000 sq. ft. Repeat as necessary. To Avoid occasional injury to cucumbers, melons and squash do not apply when plants are wet with dew. For peppers and celery, work into top few inches of soil at time of planting. Use 6 oz. per 1000 sq. ft. (15 lbs. per acre) for cutworms; for wireworm on peppers and celery, use 10 oz. per 1000 sq. ft. (25 lbs. per acre). Do not apply to any of the above plants after fruit, pods or edible parts begin to form.

For OUTSIDE HOUSEHOLD USE to control Ants, dust or spray along runways, cracks in walks and patios, around garbage cans, under porches and around outsides of house foundations. Spot treat nests in lawns and around buildings. For spiders, spray webs. For wasps, apply spray as a drench to nest under eaves, porches and sheds. For maximum kill apply in evening. Do not use in dwellings or dairy barns.

**CAUTION:** Avoid breathing dust or spray mist. Harmful if swallowed. Contact with the skin can cause toxic symptoms. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not contaminate drinking water or use near fish ponds nor birdbaths. Do not feed treated forage (pea vine) to Dairy or Beef Cattle. Dispose of container promptly when empty. **KEEP OUT OF REACH OF CHILDREN.**

This product is toxic to fish and wildlife.

U.S.D.A. Reg. No. 572-173



ACTIVE INGREDIENT: Technical Chlordane\* 10.0%  
 INERT INGREDIENTS: 90.0%

\*Equivalent to 6.0% Octachloro-4,7-Methanotetrahydroindane and 4.0% related compounds, on a weight basis.

**"HOMEOWNER" 10% CHLORDANE DUST**

**DIRECTIONS FOR USE**

To Make Spray Solution: Mix 2 tablespoons with a little water to make a thin paste. Stir paste into 1 gallon of water. Agitate while using. For Hose-End Sprayers follow manufacturers directions.

To control Ants, Leaf Miners, Leaf Rollers, Thrips, Plant Bugs, Leaf Beetles, Lace Bugs, Japanese Beetles (Adults) and certain Caterpillars on FLOWERS AND ORNAMENTALS: Dust or spray all plant surfaces in the early morning or evening when the air is still. For Taxus Weevils, treat lower branches and soil around the base of the plants. For Earwigs, Crickets and Mole Crickets use 1 pound for 700 sq. ft. of soil around plants. Repeat as necessary.

For Japanese Beetle Grubs, White Grubs and Chafer Beetle Grubs in LAWNS, apply 1 pound to 700 sq. ft. of lawn area and water thoroughly to carry insecticide into soil. For Lawn Moths, Sod Webworms, Mole Crickets and Crickets, water lightly after application. Keep children and pets off treated area until thoroughly dry. May be harmful to birds feeding on treated areas.

USE AS DUST OR SPRAY  
 FOR LAWNS AND GARDENS



Net Weight One Pound



In VEGETABLE GARDEN, for Flea Beetles and Hornworms on tomatoes, Flea Beetles on beans, black-eyed peas, eggplant and cabbage, Cucumber Beetle, Squash Bug and Squash Vine Borer on cucumbers, melons and squash, dust 4 to 6 oz.

CHLORDANE 10% DUST per 1000 sq. ft. Repeat as necessary. To Avoid occasional injury to cucumbers, melons and squash do not apply when plants are wet with dew. For peppers and celery, work into top few inches of soil at time of planting. Use 6 oz. per 1000 sq. ft. (15 lbs. per acre) for cutworms; for wireworm on peppers and celery, use 10 oz. per 1000 sq. ft. (25 lbs. per acre). Do not apply to any of the above plants after fruit, pods or edible parts begin to form.

For OUTSIDE HOUSEHOLD USE to control Ants, dust or spray along runways, cracks in walks and patios, around garbage cans, under porches and around outsides of house foundations. Spot treat nests in lawns and around buildings. For spiders, spray webs. For wasps, apply spray as a drench to nest under eaves, porches and sheds. For maximum kill apply in evening. Do not use in dwellings or dairy barns.

CAUTION: Avoid breathing dust or spray mist. Harmful if swallowed. Contact with the skin may cause toxic symptoms. In case of contact with skin, wash with soap and water. Avoid contamination of food and feedstuffs. Do not contaminate drinking water or use near fish ponds or birdbaths. Do not feed treated forage (pea vine) to Dairy or Beef Cattle. Dispose of container promptly when empty. KEEP OUT OF REACH OF CHILDREN.

This product is toxic to fish and wildlife.

© 1964, W. R. Grace & Co., Inc.