




NET CONTENTS FOUR POUNDS

ROCKLAND

**3% CIODRIN
INSECTICIDE**

LIVESTOCK DUSTING POWDER



*A Ciodrin[®] Insecticide Dust for Direct
Application to*

DAIRY AND BEEF CATTLE

*For the Control of Horn Flies—and to Lactating
Dairy Animals, Beef Cattle and Swine as an aid
in the Control of Lice.*


CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR COMPLETE CAUTION STATEMENT

ACTIVE INGREDIENTS:	BY WEIGHT
Dimethyl Phosphate of Alpha-Methylbenzyl 3-hydroxy-cis crotonate	3.0%*
INERT INGREDIENTS	97.0%
*CIODRIN [®] Insecticide, Shell Chemical Co.	TOTAL -- 100.0%

Rockland Chemical Co., Inc.

Passaic Avenue, West Caldwell, N. J.



GENERAL DIRECTIONS

ROCKLAND 3% CIODRIN[®] Insecticide is a specially prepared ready-to-use livestock dusting powder suitable for use in conventional hand and power dusters. No feed-off period is required when used on animals being fed for slaughter. Do not treat calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

USE AND DIRECTIONS FOR APPLICATION

BEEF AND LACTATING DAIRY ANIMALS:

For the control of horn flies, and as an aid in the control of lice, apply 1-2 heaping tablespoonsful (1-2 oz.) to the backline of animal covering in addition the poll, neck and upper portions of the sides. Repeat above application as necessary to maintain control, but not more often than every 14 days. Wash hands thoroughly after use.

As an aid in the control of Lice on SWINE, apply 1-2 oz. as thoroughly as possible with attention to the neck and around the ears. Repeat in 3-4 weeks if necessary.

DUST BAG-ANIMAL DUSTING STATIONS

For control of hornflies on dairy cattle and beef cattle and as an aid in the control of lice on dairy cattle, apply daily when application is made with Dust Bags. To prepare dust bags for animal use, place 4-8 pounds of 3% CIODRIN INSECTICIDE DUST in cotton cloth or double burlap bags and suspend bags near mineral or salt blocks, in alleyways leading to or from animal buildings or dairy barns. Dust bags may also be suspended in livestock holding pens, feed lots and loafing sheds where possible. Protect bags from weather by placing them under shelter.

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid inhalation of dust and contact with skin. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Keep away from open flame. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. **ANTIDOTE:** If swallowed, induce vomiting by sticking finger down throat or by giving warm salt water (1 tablespoon salt to a glass of water). **CALL A PHYSICIAN AT ONCE.** Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

DESTROY USED CONTAINERS. Do not re-use for any purpose. Destroy container by burning or burying in an isolated area. If burned, keep out of smoke.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 572-175
EPA Est. 572-NJ-1