

TREATS **2500** SQ. FT.

ROCKLAND

CRABGRASS PREVENTER "T"

WITH
TUPERSAN[®]

ROCKLAND CRABGRASS PREVENTER "T"^{with} TUPERSAN is highly effective for the control of crabgrass (smooth and hairy) and certain annual weed grasses in turf areas such as golf course fairways, lawns, parks, roadsides, and turf grown for grass seed or sod production. Because of its high degree of selectivity, CRABGRASS PREVENTER "T"^{with} TUPERSAN may be applied at time of seeding without causing injury to germinating seeds of many common temperate region grasses as well as established turf.

ROCKLAND CRABGRASS PREVENTER "T"^{with} TUPERSAN is recommended for pre-emergence control of annual weed grasses, such as crabgrass (smooth and hairy), foxtail, and barnyardgrass, in newly seeded or established plantings of the following grasses only: bluegrass, fescue, redtop, smooth brome, perennial ryegrass, orchardgrass, and the following strains of bentgrass: Penncross, Seaside, Highland, Astoria, Nimisila, C-1, C-7, and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result. Treatment will not control annual bluegrass (*Poa annua*), clover or most broadleaf weeds. CRABGRASS PREVENTER "T"^{with} TUPERSAN may be applied at any stage of turf development, including new seedings.

activity, CRABGRASS PREVENTER TUPERSAN may be applied at time of seeding without causing injury to germinating seeds of many common temperate region grasses as well as established turf.

ROCKLAND CRABGRASS PREVENTER "T" ^{with}TUPERSAN is recommended for pre-emergence control of annual weed grasses, such as crabgrass (smooth and hairy), foxtail, and barnyardgrass, in newly seeded or established plantings of the following grasses only: bluegrass, fescue, redbud, smooth brome, perennial ryegrass, orchardgrass, and the following strains of bentgrass: Penncross, Seaside, Highland, Astoria, Nimisila, C-1, C-7, and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result. Treatment will not control annual bluegrass (*Poa annua*), clover or most broadleaf weeds. CRABGRASS PREVENTER "T" ^{with}TUPERSAN may be applied at any stage of turf development, including new seedings.

DIRECTIONS

Before making application lawn should be raked to remove extraneous material and insure uniform application. Use any spreader in good condition. Measure off area to be treated. Do not spread over larger area than recommended. Water must be provided by rainfall or normal lawn watering within 3 days after application.

NEW SPRING SEEDINGS: Apply contents of this bag over 5000 sq. ft. of lawn area as final operation following seeding.


ESTABLISHED TURF AND NEW FALL SEEDINGS: Apply contents of this bag over 2500 sq. ft. of lawn area in the Spring just before expected emergence of annual weed grasses.

SUGGESTED SPREADER SETTINGS

SPREADER MODEL	NEWLY SEEDED LAWNS	ESTABLISHED AND FALL SEEDED LAWNS
SCOTT	6½	9½
MODERN	6½	9½
LAWN BEAUTY	6½	9
PATCO	6½	8¾
JACKSON	6	8¾
CYCLONE	4.5	5.8

CAUTION: Keep out of reach of children. Do not store near seeds, fertilizers and other pesticides. Do not contaminate feed or foodstuffs. Do not take internally. Avoid contact with eyes. Wash thoroughly after using. Use only on turf areas, do not apply to crops used for food and feed purposes. Store in dry place.

U. S. D. A. Reg. No. 112-179



TREATS

7500 SQUARE FEET

- ① Containing 2,4-D and special wetting agent.
- ② Kills Dandelions and Plantain
- ③ For best results apply when grass is newly mowed and moist.

CAUTION


KEEP OUT OF REACH OF CHILDREN
 SEE PAGE THREE FOR COMPLETE CAUTION
 READ ENTIRE LABEL BEFORE USING THIS PRODUCT

REGISTERED
 PATENT
 TRADE MARK
 U.S. PATENT OFFICE
 572-1827

ACTIVE INGREDIENTS
 Quantity...
 NET WEIGHT

MANUFACTURED BY NET WT 10 LBS

ROCKLAND CHEMICAL CO., INC.
 WEST CAPEWELL, N. J.



TREATS **7500** SQ. F

ROCKLAND DANDELION KILLER

DIRECTIONS

In all non-cropland apply ROCKLAND DANDELION KILLER as it comes from the bag or in any well mixed material. Measure off area to be treated.

For best results apply in a broadcast manner when grass is matted (from morning or evening dew). Apply when weeds are in active growth. On recently seeded lawn, apply only after the grasses are well established and have been mowed. Do not apply on a sidewalk to avoid drifting of material.

Use a spreader to apply to lawn, not a sprayer and apply. Do not allow to come in contact with flowers, shrubs, evergreens or other desirable plants. Be certain you are not applying to them or over that to spread.

RECOMMENDED SPREADER SETTINGS

COLE	10	PACCO	0
MOORE	0	JACKSON	0
WALKER	0	OXFORD	0

For best results apply uniformly by hand or with spreader in sufficient. The contents of this bag covers 7500 square feet. Use proportionate amounts for smaller areas (1 lb. per 750 sq. ft.). Use synthetic rubber gloves when apply.