



ROCKLAND

Net
Contents
8 ounces



- ① **Easy to use in home gardens.**
- ② **Effectively controls aphids.**
- ③ **Multi-purpose insecticide.**

**CAUTION: Keep out of reach of children.
See back panel for additional caution.**

ACTIVE INGREDIENTS:

Malathion*
Xylol

INERT INGREDIENTS:

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

DIRECTIONS FOR USE

MIXING AND APPLICATION: Mix the recommended amounts of PREMIUM MALATHION in water until the spray is uniform. Thoroughly wet plants with the spray mixture, especially the upper and undersides of the leaves. Make applications as needed except where TWO or more are indicated below.

DILUTION NOTE: 1 teaspoonful per gallon of water equivalent to 1 pint per 100 gallons of water. NOTE: The rate apply to sprayers attached to a garden hose as well as conventional equipment.

ORNAMENTALS: For the control of Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle, adult Leaf Bug, Tarnished Plant Bug, Rose Leafhopper and Potato Leaf Hopper, use 1 1/2 pints per 100 gallons of water.

For control of Soft Brown Scale, Monterey Pine Scale and Black Scale Crawlers, use 2 1/2 pints per 100 gallons.

For control of Juniper Scale, Fletcher Scale, Bagworms and Oak Kermes, use 2 pints to 100 gallons of water, crawlers have settled on the foliage.

For control of Pine Needle Scale, use 4 pints per 100 gallons of water. Apply after crawlers have settled on foliage.

NOTE: Do not use on Boston Maidenhair, Pteris Ferns as well as species of Crassula. Do not apply on Petunia.

APPLES: Use 1 to 2 pints per 100 gallons of water for the control of European Red Mite, Two Spotted Mite, Will and Clover Mite, making two or more applications as needed. For the control of Woolly Apple Aphid, Green and Rosy Apple Aphid, make one or more applications as needed. May injure McIntosh and Cortland apples if applied in 4 weeks of harvest.

PEARS: Use 1 to 2 pints per 100 gallons of water for control of Spider Mites, making two or more applications. For control of Pear Psylla, make one or more applications as needed.

TOMATO: For Spider Mites, use 1 1/2 pints to 100 gallons of water. For Aphids, use 1 pint to 100 gallons of water. Applications later than 10 days before harvest.

PEAS: For Pea Aphids, use 1 1/2 pints to 100 gallons of water. Make no applications later than 14 days before harvest.

BEANS: For Spider Mites, use 1 to 1 1/2 pints to 100 gallons of water. For Mexican Bean Beetle, use 1 1/2 pints to 100 gallons of water. Make TWO or more applications as needed. Make no applications later than 10 days before harvest.

NOTE: Make no application on APPLES and PEARS later than 14 days before harvest.

CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist or dust. Avoid repeated or prolonged contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children away from heat or flame.

This product is toxic to fish. Keep out of lakes, streams and ponds.

NOTE TO PHYSICIAN: Malathion, upon repeated, prolonged and/or careless use, may cause cholinesterase inhibition. Atropine is antidotal.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to the instructions.

U.S.D.A. Reg.

ROCKLAND CHEMICAL CO., Inc.
Passaic Avenue West Caldwell, N. J.

ACTIVE INGREDIENTS:

Malathion*

Xylol

INERT INGREDIENTS:

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

50.0%

40.8%

9.2%

DIRECTIONS FOR USE

MIXING AND APPLICATION: Mix the recommended amounts of PREMIUM MALATHION in water until the spray mixture is uniform. Thoroughly wet plants with the spray mixture, especially the upper and undersides of the leaves. Make one or more applications as needed except where TWO or more are indicated below.

DILUTION NOTE: 1 teaspoonful per gal. of water equivalent to 1 pint per 100 gals. of water. **NOTE:** The rate of dilutions apply to sprayers attached to a garden hose as well as conventional equipment.

ORNAMENTALS: For the control of Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle adults, Fourlined Leaf Bug, Tarnished Plant Bug, Rose Leafhopper and Potato Leaf Hopper, use 1 1/2 pints per 100 gallons of water.

For control of: Soft Brown Scale, Monterey Pine Scale and Black Scale Crawlers, use 2 1/2 pints per 100 gallons of water.

For control of: Juniper Scale, Fletcher Scale, Bagworms and Oak Kermes, use 2 pints to 100 gallons of water. Apply after crawlers have settled on the foliage.

For control of: Pine Needle Scale, use 4 pints per 100 gallons of water. Apply after crawlers have settled on the foliage.

NOTE: Do not use on Boston, Maidenhair, Pteris Ferns as well as species of Crassula. Do not apply on Petunias.

APPLES: Use 1 to 2 pints per 100 gallons of water for the control of: European Red Mite, Two-Spotted Mite, Williamette Mite and Clover Mite making two or more applications as needed. For the control of Woolly Apple Aphid, Green Apple Aphid and Rosy Apple Aphid, make one or more applications as needed. May injure McIntosh and Cortland apples if applied within 4 weeks of harvest.

PEARS: Use 1 to 2 pints per 100 gallons of water for control of: Spider Mites, making two or more applications as needed. For control of Pear Psylla make one or more applications as needed.

TOMATO: For Spider Mites, use 1 1/2 pints to 100 gallons of water. For Aphids, use 1 pint to 100 gallons of water. Make no applications later than 10 days before harvest.

PEAS: For Pea Aphids, use 1 1/2 pints to 100 gallons of water. Make no applications later than 14 days before harvest.

BEANS: For Spider Mites, use 1 to 1 1/2 pints to 100 gallons of water. For Mexican Bean Beetle use 1 1/2 pints to 100 gallons of water. Make TWO or more applications as needed. Make no applications later than 10 days before harvest.

NOTE: Make no application on APPLES and PEARS later than 14 days before harvest

CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist or dust. Avoid repeated or prolonged contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Keep away from heat or flame.

This product is toxic to fish. Keep out of lakes, streams and ponds

NOTE TO PHYSICIAN: Malathion, upon repeated, prolonged and/or careless use may cause cholinesterase inhibition. Atropine is antidotal.

WARRANTY: Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

U.S.D.A. Reg. No. 572-183

ROCKLAND CHEMICAL CO., Inc.
Passaic Avenue West Caldwell, N. J.