

ROCKLAND

NET WEIGHT 3 POUNDS

GENERAL PURPOSE DUST

FOR FRUIT TREES, SHRUBS, VEGETABLES.

CONTROLS INSECTS. PREVENTS CERTAIN DISEASES.

COMBINATION OF MALATHION, METHOXYCHLOR AND CAPTAN

CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR COMPLETE CAUTION STATEMENT.

ACTIVE INGREDIENTS:

*Malathion	5.00%
**Methoxychlor	5.00%
Captan (N-Trichloroethyl-3,4-dihydro-2H-pyridin-2-one-2,2-dicarboximide)	7.00%
INERT INGREDIENTS	83.00%

*Malathion is a registered trademark of the Cyanamid Company.
**Methoxychlor is a registered trademark of the Cyanamid Company.
Captan is a registered trademark of the Cyanamid Company.

Make dust applications on listed Flowers, Shrubs, Trees, Fruit Trees and Vegetables before insects become numerous and before diseases become apparent. Protect plants through the growing season by dusting at 7-14 day intervals and after heavy rains. Cover foliage thoroughly but do not use in excess. Pay particular attention to cover underside of leaves.

ROCKLAND GENERAL PURPOSE DUST is effective for control of the following insects and diseases:

FRUIT TREES: Aphids, Coddling Moth, European Red Mites, Clover Mites, Two Spotted Mites, Willamette Mites, Pear Psylla, Japanese Beetles, Black Cherry Aphid, Tent Caterpillar on **APPLES, PEARS, PEACHES AND CHERRIES.** Scab, Curculio on **APPLES.** Leaf Spot on **APPLES.** Brown Rot, Curculio, Scab on **PEACHES.** Curculio, Leaf Spot, Brown Rot on **CHERRIES.** Apply material directly to the tree when the air is still. For timing of applications, follow fruit tree dust schedule.

SUGGESTED FRUIT DUSTING SCHEDULE

DELAYED DORMANT — (From first green on bud to 1/2 inch leaves) on apple and peach.

PRE-BLOOM — (From first pink to first bloom) on apple, peach, pear and cherry.

PETAL-FALL — (When 90% of petals have fallen on all fruits listed above).

SHUCK-FALL — (When half the shucks have fallen from fruit) on cherry and peach.

COVER-SPRAY — (2 to 6 applications.) Begin 10 days after petal-fall on apple and pear. Begin 10 days after shuck-fall on cherry and peach. Do not apply to peaches within 21 days of harvest nor to other fruit within 14 days of harvest.

SHRUBS: Japanese Beetles, Aphids, Spider Mites, Lacebugs, Azalea Scale, Scurfy Scale, Euonymus Scale, Soft Brown Scale, Bagworms and Boxwood Leaf Miners on **AZALEAS, RHODODENDRONS, ARBORVITAE, JUNIPERS, HYDRANGEAS, EUONYMUS, PIERIS JAPONICA, LILAC AND BOXWOOD.**

NOTE: Do not use on Boston, Maidenhair, Pteris Ferns as well as species of Crassula.

TREES: Spider Mites, Birch Leaf Miners, Holly Leaf Miners, European Pine Shoot Moth, Tent Caterpillars and Aphids on **PINE, BIRCHES, HOLLY OAKS, MAGNOLIAS, MAPLES AND SYCAMORE.**

FLOWERS: Aphids, Spider Mites, Japanese Beetles, Rose Chafer, Rose Leafhoppers, Leaf Rollers, Rose Slugs, Flea Beetles, Thrips, Tarnished Plant Bugs, Four Lined Leaf Bugs, Whitefly, Asiatic Garden Beetles and Black Blister Beetles on **ROSES, GLADIOLUS, DAHLIAS, SWEET PEAS, ASTERS AND CHRYSANTHEMUMS.** Black Spot, Powdery Mildew,

SHRUBS: Japanese Beetles, Aphids, Spider Mites, Lacebugs, Azalea Scale, Scurfy Scale, Euonymus Scale, Soft Brown Scale, Bagworms and Boxwood Leaf Miners on **AZALEAS, RHODODENDRONS, ARBORVITAE, JUNIPERS, HYDRANGEAS, EUONYMUS, PIERIS JAPONICA, LILAC AND BOXWOOD.**

NOTE: Do not use on Boston, Maidenhair, Pteris Ferns as well as species of Crassula.

TREES: Spider Mites, Birch Leaf Miners, Holly Leaf Miners, European Pine Shoot Moth, Tent Caterpillars and Aphids on **PINE, BIRCHES, HOLLY OAKS, MAGNOLIAS, MAPLES AND SYCAMORE.**

FLOWERS: Aphids, Spider Mites, Japanese Beetles, Rose Chafer, Rose Leafhoppers, Leaf Rollers, Rose Slugs, Flea Beetles, Thrips, Tarnished Plant Bugs, Four Lined Leaf Bugs, Whitefly, Asiatic Garden Beetles and Black Blister Beetles on **ROSES, GLADIOLUS, DAHLIAS, SWEET PEAS, ASTERS AND CHRYSANTHEMUMS.** Black Spot, Powdery Mildew, Rust, Leaf Spot and other flower spots on **ROSES, TUBEROUS BEGONIAS, CARNATIONS, CHRYSANTHEMUMS and PEONIES.** Do not apply on Petunias.

VEGETABLES: Fruitworms, Hornworms, Aphids, Spider Mites, Flea Beetles, Colorado Potato Beetles, Early and Late Blight, Anthracnose and Septoria Leaf Spot on **TOMATOES.** Aphids, Flea Beetles, Colorado Potato Beetles and Lacebugs on **PEPPERS and EGGPLANTS.** Aphids Colorado Potato Beetles, Flea Beetles, Leafhoppers, Early and Late Blight on **POTATOES.** Pea Aphid on **PEAS**

NOTE: Make no applications on **PEAS** 14 days before harvest. Make no application on **POTATOES** 10 days before harvest. Make no applications on **TOMATOES, PEPPERS, EGGPLANTS** within 7 days of harvest.

After the time limits set above have passed, an effective fungicide should be used alone, in a manner which would not leave residues in excess of established tolerances.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing dust or spraying mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after using. Do not contaminate feed or footstuffs. Keep out of reach of children. Dispose of empty container. Do not reuse.

This product is toxic to fish. Keep out of any body of water.
WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.