



572-186
PM 12 1/4

Rockland. General Purpose Spray



ACCEPTED
SEP 22 1986
Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered in the EPA Reg. No.



****N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide**

EPA Reg. No. 572-186

EPA Est. No. 572-NJ-1

KEEP OUT OF REACH OF CHILDREN (12 pt. type)

DANGER (18 Pt. type)

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS:

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

See back panel for additional precautionary statements.

NET WEIGHT ONE POUND

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call physician or poison control center immediately. Drink 1 or two glasses of water and induce vomiting by touching finger to back of throat. Do not give anything by mouth to an unconscious person. NOTE TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal.

IF ON SKIN: Wash with plenty of soap and water. Launder clothes before reuse.

IF IN EYES: Call physician immediately, may cause irreversible eye damage. Flush eyes with plenty of water.


ENDANGERED SPECIES RESTRICTIONS

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered specie and the general location of their habitat have been identified by county:

Freshwater Mussels: Virginia (Smyth, Scott, Lee, Washington and Grayson)
Prior to the use of this product in the above counties, you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service office (Endangered Species Specialist) or State Wildlife Agency if there is any doubt as to the areas where the outdoor use of captan is prohibited.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.



Wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan when using indoors or outdoors. When using outdoors wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

Make spray application on listed Flowers, Shrubs, Trees, Fruit Trees and Vegetables before insects become numerous and before diseases become apparent. Mix 8 level tablespoons per gallon of water (1 lb. per 8 gallons of water). Mix thoroughly before using and occasionally during use to assure uniformity of spray mix. For hose-end sprayer, follow manufacturer's directions. Protect plants through the growing season by spraying at 7-14 day intervals and after heavy rains. Do not store diluted spray. Pay particular attention to cover underside of leaves.

ROCKLAND GENERAL PURPOSE SPRAY is effective for control of the following insects and diseases:

FRUIT TREES: Aphids, Codling Moth, European Red Mites, Clover Mites, Two Spotted Mites, Willamette Mites, Pear Psylla, Japanese Beetles, Black Cherry Aphid, Tent Caterpillar on APPLES, PEARS, PEACHES AND CHERRIES, Scab, Curculio on APPLES, Leaf Spot on PEARS, Brown Rot, Curculio, Scab on PEACHES, Curculio Leaf Spot, Brown Rot on CHERRIES. Apply material directly to the tree when the air is still. For timing of applications, follow fruit tree dust schedule.

SUGGESTED FRUIT SPRAYING SCHEDULE

DELAYED DORMANT - (From first green on bud to 1/2 inch leaves) on apple and peach.

PRE-BLOOM - (From first pink to first bloom) on apple, peach, pear and cherry.

PETAL-FALL - (When 90% of petals have fallen on all fruits listed above).

SHUCK-FALL - (When half the shucks have fallen from fruit) on cherry and peach.

COVER-SPRAY - (2 to 8 applications.) Begin 10 days after petal fall on apple and pear. Begin 10 days after shuck-fall on cherry and peach. Do not apply to peaches within 21 days of harvest nor to other fruit within 14 days of harvest.

SHRUBS: Japanese Beetles, Aphids, Spider Mites, Lacebugs, Azalea Scale, Scurfy Scale, Euonymus Scale, Soft Brown Scale, Bagworms and Boxwood Leaf Miners on AZALEAS, RHODODENDRONS, ARBORVITAE, JUNPERS, HYDRANGEAS, EUONYMUS, PIERIS JAPONICA, LILAC AND BOXWOOD. NOTE: For Scale insects and Bagworms on SHRUBS, make ONLY dust applications. Do not spray for control of these insects as 2 to 3 times the normal dosage of this product is required per gallon of water. This would result in excessive clogging of spray nozzles.

NOTE: Do not use on Boston, Moldenhair, Pieris Ferns as well as species of Crassula.

TREES: Spider Mites, Birch Leaf Miners, Holly Leaf Miners, European Pine Shoot Moth, Tent Caterpillars and Aphids on PINE, BIRCHES, HOLLY OAKS, MAGNOLIAS, MAPLES AND SYCAMORE.

FLOWERS: Aphids, Spider Mites, Japanese Beetles, Rose Chafer, Rose Leafhoppers, Leaf Rollers, Rose Slugs, Flea Beetles, Thrips, Tarnished Plant Bugs, Four Lined Leaf Bugs, Whitefly, Asiatic Garden Beetles and Black Blister Beetles on ROSES, GLADIOLUS, DAHLIAS, SWEET PEAS, ASTERS AND CHRYSANTHEMUMS. Black Spot, Powdery Mildew, Rust, Leaf Spot and Botrytis Flower Blight on ROSES, TUBEROUS BEGONIAS, CARNATIONS, CHRYSANTHEMUMS and PEONIES. Do not apply on Petunias.

VEGETABLES: Do not rotate a foliar treated crop with crops other than those with registered captan uses.

... Fruitworms, Hornworms, Aphids, Spider Mites, Flea Beetles, Colorado Potato Beetles, Early and Late Blight, Anthracnose and Septoria Leaf Spot on TOMATOES, Aphids, Flea Beetles, Colorado Potato Beetles and Lacebugs on PEPPER and EGGPLANTS, Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers, Early and Late Blight on POTATOES, Pear Aphid on PEAS.

NOTE: Make no applications on PEAS 14 days before harvest. Make no application on POTATOES within 10 days before harvest. Make no applications on TOMATOES, PEPPERS, EGGPLANTS within 7 days of harvest.

After the time limits set above have passed, an effective fungicide should be used alone, in a manner which would not leave residues in excess of established tolerances.

DO NOT ALLOW PERSONS TO ENTER TREATED AREAS WITHIN 4 DAYS FOLLOWING APPLICATION UNLESS PROTECTIVE CLOTHING IS WORN. Conspicuously post reentry information at site of application.

STORAGE: Store unused material in original container only, in a locked, dry storage area out of reach of children and animals. Keep container tightly closed. Do not contaminate water, food or feed by storage or disposal.

DISPOSAL: Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by



Rockland Chemical Co., Inc.
West Caldwell, NJ 07006 U.S.A.

© Rockland Chemical Co., Inc. 1983

Product A18601

BEST AVAILABLE COPY