

ROCKLAND WEED KILLER is highly effective as a contact spray for control of weeds and grasses and for killing undesirable woody plants. Temporary non-productivity of soil may be caused by heavy applications of the product. This condition usually disappears in the over-winter period. Some species of woody plants and perennial weeds are difficult to control and re-treatment may be necessary if re-growth occurs.


This product is non-volatile and non-flammable.

WEED AND GRASS CONTROL

For control of: Woody Perennials such as poison ivy, poison oak, and poison sumac; Perennials such as leafy spurge, bitter dock, goldenrod, perennial ragweed milkweed, and blueweed; and Annuals such as crabgrass, broomweed, chickweed, cocklebur imson weed, lambsquarters, larkspur prickly lettuce, ragweed and shepherdspurse.

Mix I gallon ROCKLAND WEED KILLER with 5 to 7 gallons of water (I quart with 5 to 7 quarts of water). Thoroughly wet foliage and stems of undesired vegetation. The degree of control and duration of effect will vary with weed species, rainfall, temperature and other conditions.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use or this product other than indicated on the label Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions



BRUSH CONTROL

Rights of Way - Roadsides Drainage Ditches - Rangelands - Pastures

ROCKLAND WEED KILLER applied as a foliar spray, is highly effective for killing most woody plants, including hardwood and coniferous species such as alder, ash, bitch, cedar, elm, gum, hickory, maple, oak, pine, and willow.

Prepare spray solution by mixing one gallon with 5 to 7 gallons of water.

Spray any time after brush has reached the full-leaf stage until foliage begins to discolor, preferably during period of high temperatures and high humidity. Apply as a full-coverage spray to foliage, stems, limbs, and base of brush; thorough coverage is essential for best results. In roadside applications on tall, dense brush, it is often necessary to spray from field side as well as road side. Spraying away from crops planted close to rights of way will aid in preventing contact of crops with spray or drift; coarse sprays are less likely to drift. The product is not volatile.

CAUTION: Avoid prolonged contact with skin. In case of contact, wash off with plenty of water.

Do not apply to, or drain, or flush equipment on desirable plants or vegetation, as injury may result. Do not drain or flush equipment near domestic waters. Wash sprayer thoroughly after use to remove all herbicidal solution and to reduce corrosion.

U.S.D.A. Reg. No. 572 212

ROCKLAND

TREATS 10,000 SQ. FT.

PROFESSIONAL CRABGRASS PREVENTER WITH BALAN* ACCEPTED

For Professional Use Only

A Balan-coated corn cob for pre-emergence control of crabgrass and annual blue grass in established lawns.

DIRECTIONS FOR USE

LAWN and ORNAMENTAL TURFS (established turf only):

Apply contents of bag (25 lbs.) to 10,000 sq. ft. (100' x 100') o a spreader for the control of SMOOTH and HAIRY CRABGRASS, GOOSEGRASS (Line): Crabgrass), BARNYARD-GRASS, YELLOW and GREEN FOXTAIL and ANNUAL BLUEGRASS (Poa Annua). This product controls weeds by killing their seeds as they germinate. It does not control established weeds. Time application in the Spring before the germination of annual grass weeds listed. Water after application to wash chemical into the turf.

SPREADER SETTING: CYCLONE 3

PRECAUTION: Do not apply to turf planted the previous Fall. Do not apply to bentgrass putting greens or dichondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustine-grass or Centipedegrass until these turfs are well-established since root inhibition may result. Do not reseed within 42 days after treatment.

ACTIVE INGREDIENT:

N-butyl Niethyl a a a-trifucro-2 6 dicitro folulaine 1.9
INERT INGREDIENTS: 98.1
*BALAN—the registered from many for benefic Europ Products Co

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or absorbed through the skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. In case of contact flush with water. Wash thoroughly with soap and water after use. Do not contaminate foodstuffs or feed.

Do not feed forage from treated areas to livestock. Destroy bag when empty. Do not reuse.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application cleaning of equipment or disposal of wastes.


Net Weight 25 lbs.

Manufactured by

ROCKLAND CHEMICAL CO.

PASSAIC AVENUE

WEST CALDWELL, NEW JERSEY



the was the same and the same

The state of the state of the state of the state of

The same of the sa



TUPERSAM

TREATS

5000 SQUARE FE

The Mark Stroken AM

- (1) Controls Crabgrass and other Annual Weeds and Grasses in Turf.
- threeds with 18-6-6 Fertilize
- Permits seeding when used as directed.

READ ENTIR

CAUTION: KEEP OUT FACH OF CHICARIA

SEE BACK FOR COMPLETE SYSTEM FERRENCE NO. 1

Trailing a phase through the con-

MANUFACTURES BY

THE TON SE

ROCKLAND CHEMICAL CO., INC.

WEST CALDWELL, N. J. 07006