SEST AVAILABLE COPY

INSECT SPRAY AND TERMITE PROOFER

CONTROLS: CHINCH BUGS . CERTAIN WHITE GRUB . HYPERODES WEEVILS . BAGWORMS . WOOD BORERS . AND OTHER INSECTS AS LISTED.

RECOMMENDED FOR AREA CONTROL OF SUBTERRANEAN TERMITES

ACTIVE INGREDIENTS

Chlorpyrifos [0,0-diethyl 0-(3, 5, 6-trichloro-2- pyridyl) phosphorothicate].....13.5% Aromatic Petroleum Solvent......81.5% INERT INGREDIENTS.....5.0%

1 ---100.0% Contains 1 pound of Dursban* (chlorpyrifos) per gallon. *DURSBANR Trademark of the Dow Chemcal Co. U.S. Patent No. 3,244,586.

KEEP OUT OF REACH OF CHILDREN - (12 pt type)

WARNING - (18 pt. type)

See Back Panel for Additional Precautionary Statements

 Ur^{-} Fr. **6**. .

STATEMENT OF PRACTICAL TREATMENT

P+ -:

Call a physician or nearest Poison Control Center if a Known exposure occurs or if poisoning is suspected.

IF SWALLOWED: D not induce vomiting. Contains Aromatic Petroleum Solvent. Call a physician. IF ON SKIN: Remove contaminated clothing and immediately wash skin with soap and water. Launder clothes before reusing. IF IN EYES: Flush with plenty of water, get medical attention. IF INHALED: Remove inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

EPA REG. NO. 572-213

EST. NO. 572-NU-1

NET CONTENTS:

Back Panel

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: WARNING-May be fatal if swallowed. May be absorbed through the skir. To not breath vapors or spray mist. Do not get on skin, eyes or cloching. Wash thoroughly after handling. Handle concentrate in ventilated area. Keep pets and child: in off treated areas until dry.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and other wildlife. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply where runoff is likely to occur or when weather conditions favor drift from treated area. Cover bird baths and small fish pools. This product is highly toxic to bees exposed to direct treatment or residue on plants. Protective information may be obtained from our Cooperative Agricultural Extension Service.

Mail A COPY

Hell.

ROCKLAND INSECT SPRAY AND TERMITE PROOFER dilutes in water to make a contact and residual spray for the control of many turf and ornamental insects.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PESTS OF LAWNS AND OTHER ORNAMENTAL AND RECREATIONAL TURFGRASS AREAS: Use ROCKLAND INSECT SPRAY AND TERMITE PROOFER to control ants, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers and dos webworms (lawn-moths). Mix with water according to directions given in the following table and apply as a coarse, low-pressure spray using suitable application equipment. For best results, the lawn should be moist at time of treatment and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; repeat when needed. Keep children and pets off treated areas ur il spray has dried. Do not graze treated areas.

DOSAGE and DILUTIONS for USE on TURF

reas to be Treated	AMOUNT OF ROCKLAND INSECT SPRAY AND TERMITE PROOFER	Diluted with Water To Give
00 sq. ft. 00 sq. ft. 00 sq. ft. 700 sq. ft. One Acre	1.5 fl. oz. 3.0 fl. oz. 15.0 fl. oz. 2.0 pints 4.0 pints (2 tablespoors l fl. oz.)	15 gallons 30 gallons 150 gallons 325 gallons 1,300 gallons

To control certain white grubs (larvae of European chafer and Japanese beetle).

Use two to four times the amounts of ROCKLANI INSECT SPRAY AND TERMITE PROOFER shown in the TURF table, i.e., 6 to 12 fl. oz. per 1,000 sq. ft. or 2 to 4 sallons per acre. Apply in enough water to obtain uniform coverage of the area followed immediately by thorough watering to wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

To control Turfgrass (Hyperodes) weevils: Use twice the amount of ROCKLAND INSECT SPRAY AND TERMITE PROOFER shown in TURF table, i.e., 2 gallons diluted in 1,300 gallons of water per acre. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

MOTE: To help prevent infestation of buildings by ants, crickets, earwigs, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and diacent to the building, also the building foundation to a height of 2 to 3 ft; there pests are active and may find entrance. Use the same dosage and dilution' commended for control of these pests on turf as shown in the TURF table. Apply a coarse spray to thoroughly and uniformly wet the band area.

BEST AVAILABLE COPT

NOTE: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for volving periods of time after treatment, depending on exposure of treated reas to weathering conditions.

PESTS OF ORNAMENTALS: Use ROCKLAND INSECT SPRAY AND TERMINTE PROOFER to treat flowers, shrubs, vines, shade and flowering trees, and evergreens found to be infested with pests listed in the following table. Dilute with water according to directions. Apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage. For best results, apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes, or variegated ivy because of possible injury to these plants.

Dilute the directed amount of ROCKLAND INSECT SPRAY AND TERMITE PROOFER with water to make the desired amount of spray mix. 1 fl. oz. to 3 gallons--3.2 fl. o to 10 gallons--1 quart to 100 gallons to control Aphids, Bagworms, Eastern Tent Caterpillars, Fall Webworm, Grasshoppers, Mites, Orange Stripped Oakworm, Whiteflies, Yellow-necked caterpilla, Gypsy Moth. Use 2 fl. oz. to 3 gallons--6.4 fl. oz. to 10 gallons--2 quarts to 100 gallons to control Cutworms, Leafhoppers, Mealybugs, Mimose Webworm, Redhumped Caterpillar, Thrips. Use 4 fl. oz. to 3 gallons--13 fl. oz. to 10 gallons--1 gallon to 100 gallons for WOOD BORING INSECTS infesting such trees and shrubs as Dogwood, Lilac, Ash, Rhododendron and Apply as in a coarse, wet spray to the trunks and lower limbs when adults begin to emerge. Some wood borers require several applications. Timing of sprays is critical to best control. Consult your State Cooperative Agricultur Extension Service for proper timing in your location. For Peach Tree Borer on flowering trees and shrubs of the Genus Prunus apply as a trunk spray before newly hatched larvae enter the tree. Thoroughly wet all bark areas from the grou to scaffold limbs. Consult your Extension Service specialist for proper timing.

SUBTERRANEAN TERMITES-Directions for Homeowner use.

DILUTIONS TO USE ON TERMITES

ROCKLAND INSECT SPRAY AND TERMITE PROOFER	Water =	Linear Ft. Treated
16 oz. 1 qt. 1 gal.	6 gml.	8 Lin. Ft. 16 Lin. Ft. 64 Lin. Ft.

TERMITE CONTROL IN EXISTING STRUCTURES -BUILDINGS WITH BASEMENTS OR CRAWL-SE'ACES

First remove all wood debrie and wood forms. Around piers, pipes, chimney bases and along foundations of poured concrete dig a trench 6 inches wide and inches deep. For brick and block foundations dig the trench about 12 inches deep. If footing is more than 12 inches deep, make holes with a crowbar, pipe for rod about one foot apart that extend from the trench bottom to the top of the footing. Make holes closer in hard-pa i clay soils. NEVER DIG BELGH THE TOP OF THE POOTING. Apply the emulsion (o. olution) in a sprinkler can in the trench at a rate of 4 gallons per 10 linear feet per foot of depth. Apply half of this to the back-fill. Likewise, treat along the inside of foundations of crawl space buildings. Cover treated soil with a thin layer of uncreated soil.



For raised porches, terraces, and entrance slabs, drill holes at one foot intervals, for inches from the foundation, and pour one-half gallon of emulsion (or solution) in inches from the foundation, and pour one-half gallon of emulsion (or solution) in inches from the foundation of the foundation of

The object is to extablish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment, particularly if there are inaccessible areas, basements or slab-type construction. It is suggested that you contact the Agricultural Cooperative Extension Service for advice or consult a Commercial Pest Control Service.

Retreatment of subterranean termites should only be made when there is evidence of reinfestation subsequent to the initial treatment, or there has been a distruption of the chemical barrier in the soil due to construction, excavations, landscaping, etc. Reapplication should be made as a spot treatment to these areas.

Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the termite chemical solution while filling or mixing this material with water. Use anit-back-flow or anti-siphonage equipment on all filling equipment. These devices can be purchased at hardware, or plumbing supply stores. Use of antiback-flow equipment will help to insure that the user of this toxic chemical will mot contaminate domestic water supplies. Refer to Federal (Federal Housing Administration), state, and local specifications for safe distances of treatment areas from wells. Soil in the vicinity of wells, cisterns or ponds should not be treated if it is water saturated, subject to excessive saturation due to an accumulation of water or if the soil to be treated around the perimeter of the structure is at a grade lower than surrounding yard ; property area. If this is the situation, treat as follows: Do Not Apply Under hardsure. Soil should be removed to an area safe from well or domestic water contamination, theated, allowed to stand undisturbed for two to four hours then returned to the trench which has been lined with 4 mil. plastic sheeting. Se careful not to puncture the plastic sheeting when returning soil to the trench. Do not treat structures that contain cisterns or wells.

STORAGE AND DISPOSAL

STORAGE: Store unused material, in original container only, in a locked storage area out of reach of children and animals. Keep container tightly closed. Do not contaminate water, food or feed by storage or disposal. Do not use, pour, spill or store near heat or open flame.

PESTICEDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

CONTAINER DISPOSAL: UP TO ONE GALLON: Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash. Rinse thoroughly, before discarding in trash. FIVE GALLONS & LARGER - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or, puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities,

WARRANTY: Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA REG No. 572-213

EST No. 572-NJ-1

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WN-567) WASHINGTON, D.C. 20440

7056-1 23

NOTICE OF PESTICIDE: PAEGISTRATION

REBEGIST TATION

(Under the Federal Insecticide, Fundic to and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

CSA Surface Spray Formula Two

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

CSA Limited, Inc. P.O. PCX 73308 Houston, TX 77273

BEST AVAILABLE COPY

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the libeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pristicide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the regist ant a right to exclusive use of the name or to its use if it has been covered by others.

This product is condit. ally registered in accordance with FIFFA sec. 3(c)(7)(8) provided that v

- 5. Submit and/or cite ... data required for registration/reregistration. of your product under FIRS one, 3(c)(5) when the Agency requires all registrants of similar reads to submit such data.
 - Make the labeling riges listed below before you release the product or ab iment:
 - Id the phrase "FFF Temperation No. 7056-123."
- Perise your "Fest; " Disposal" directions to read:

Wastes result it from the use of this product may be disposed of on sit < 'ar approved waste disposal facility.

3. Submit five (5) corres of your final printed labeling before you release the product for element. Refer to the A-79 Enclosure for a further description of final print blabeling.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

1/1/1

EPA Firm \$570-6 (Rev. 5-76) 📦

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.