


ACCEPTED
MAY 1 1939
UNDER FEDERAL INSECTICIDE
ACT AND FEDERAL FUNGICIDE ACT
REGISTERED UNDER NO. 572-214



ROCKLAND

\$5.95



**LAWN
FUNGICIDE
WITH DYRENE***

**TREATS
4000 SQ. FT. PREVENTIVE**

TREATS
4000 SQ. FT. PREVENTIVE

- Controls and Prevents Lawn Diseases.
- Apply for Brown Patch, Melting Out, Snow Mold, Leaf Spot and others listed.
- Do not water or mow 24 hours after application.

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR COMPLETE CAUTION
READ ENTIRE LABEL BEFORE USING THIS PRODUCT

ACTIVE INGREDIENT:
2, 4-Dichloro-6-g-chloroanilino-g-triazine** — 5%
INERT INGREDIENTS: _____ 95%
100%

**U.S. Patent No. 2,728,480 and
Canadian Patent No. 544,827
Effective for purpose claimed.
©TM Chemagro Corp.


MANUFACTURED BY

NET WT. 10 LBS.

ROCKLAND CHEMICAL CO., INC.


WIE S P C / A I L D W I E 0 1 7 (0) 0 1 6

BB-18



ROCKLAND

LAWN FUNGICIDE WITH DYRENE*



ROCKLAND LAWN FUNGICIDE With DYRENE gives effective control of Rhizoctonia brown patch, Sclerotinia dollar spot, copper spot, rust, snow mold (Typhula), and Helminthosporium diseases, such as melting-out of Kentucky bluegrass, red leaf spot of bentgrasses, leaf blotch of Bermuda grass, Helminthosporium blight of fescues, brown blight of rye grasses, zonate eye spot, and Helminthosporium leaf spot of bluegrasses, bentgrasses, and fescues.

Uniform distribution of this granular material is necessary for maximum control. The ideal time for application is in the early morning when dew is on the grass, or after area has been watered, in order for granulars to adhere to the grass blades. Do not mow or water treated lawn area for 24 hours.

APPLICATION RATES

Preventive

before diseases are present.
One bag per 4000 sq. ft. lawn area

Curative

when disease damage is visible
One bag per 2500 sq. ft. lawn area

SPREADER SETTINGS

	Preventive 4000 sq. ft.	Curative 2500 sq. ft.
Jackson	6½	8½
Lawn Beauty	6½	9
Scott	7	9½
Patco	7	8½

WHEN TO APPLY

For Foliar Turfgrass Diseases listed above, begin preventive applications in the early Spring before diseases appear, and repeat at 10 to 14 day intervals. During period of wet weather applications, every 7 days may be required. Continue applications throughout the Spring and Summer months.

SPREADER SETTINGS

	Preventive 4000 sq. ft.	Curative 2500 sq. ft.
Jackson	1 1/2	1 1/2
Lawn Beauty	1 1/2	1 1/2
Scott	1 1/2	1 1/2
Patco	1 1/2	1 1/2

WHEN TO APPLY

For Follar Turfgrass Diseases listed above, begin preventive applications in the early Spring before diseases appear, and repeat at 10 to 14 day intervals. During periods of wet weather applications, every 7 days may be required. Continue applications throughout the Spring and Summer months.

NOTE: Watch for Leaf Spot in early Spring, Dollar Spot in late Spring, Brown Patch in hot, humid Summer months, and Dollar Spot and Copper Spot again in the Fall.

For Control of Snow Mold: Apply after the first killing frost in the Fall. A second application should be made when the lawn or other turfgrass area is free of snow in mid-winter. Apply a third application after the final melting of snow in late Winter or early Spring.


CAUTION

May be harmful if swallowed or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food. Keep out of reach of children.

WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

U.S.D.A. Reg. No. 572-214



W-508

