



ROCKLAND SUPER Kleen-Kow

CONCENTRATE 20-5

Ciodrin® and Vapona®

DAIRY and LIVESTOCK SPRAY



REG. U. S. PAT. OFF.

DANGER: KEEP OUT OF REACH OF CHILDREN.

See back panel for antidote statement and other warnings.

POISON




ACTIVE INGREDIENTS:

Dimethyl phosphate of alpha-methylbenzyl 3-hydroxy-cis-crotonate	20.0%
2,2-dichlorovinyl dimethyl phosphate*	4.6%
Related Compounds	.4%
Petroleum Hydrocarbons	62.0%
INERT INGREDIENTS:	13.0%

*Equivalent to 5.0% VAPONA® Insecticide
 ®VAPONA reg. trademark of Shell Chemical Co.
 ®CIODRIN reg. trademark of Shell Chemical Co.

Manufactured by

Rockland Chemical Co., Inc.
 Passaic Avenue, West Caldwell, N. J.



POISON



NET CONTENTS 1 QUART

DIRECTIONS FOR USE

For Control of Stable Flies, Horn Flies, House Flies, Lice and Lone Star Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs.

Mix 1 quart per 6 gallons of water. Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 pints of spray per large animal and proportionately less for smaller animals.

For pen or corral operations involving spray application to livestock with high pressure equipment, do not exceed a 0.5% concentration. Mix 1 quart per 12 gallons of water (0.5% concentration). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1-2 quarts of spray per large animal and proportionately less for smaller animals. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Improved Face Fly control may be achieved by adding sugar to the spray solution at the rate of 1 lb. per 6-10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Low Volume Application for Control of Face Flies

Mix 2 tablespoons (1 fluid oz.) per 1 1/2 pints of water (1% concentration). Using a calibrated sprayer, apply 5 teaspoons of spray to the face and a total of 5-10 teaspoons to the back and sides of the animal 3 times per week. Improved control may be achieved by the addition of 1 level teaspoon of sugar per pint of spray solution.

For Control of Lice and Winter Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs.

Mix 1 quart per 26 gallons of water (0.25% concentration). Spray animals thoroughly using up to 1 gallon of spray. Apply a second application 14 days later. Repeat applications as necessary to maintain control, but not oftener than once every 7 days.

Back Rubber Applications

For use in back rubbers or face rubbers for the control of Horn Flies and Face Flies mix 1 pint per 4 gallons of back rubber base oil (1% concentration). This may be diesel fuel or fuel oil in accordance with local recommendations.

For Control of House Flies, Face Flies, and Stable Flies in and around Livestock Buildings.

Mix 1 quart per 6 gallons of water (1% concentration) Spray premise, such as inside and outside barn walls (including dairy barns), eaves, fences and other resting or breeding places, at least twice monthly using approximately 1 gallon of finished

spray per 1,000 sq. ft. (40' x 25'). Improved House Fly and Face Fly control may be achieved by adding sugar to the spray solution at the rate of 1 lb. per 6-10 gallons of spray.

DANGER.

Poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Avoid prolonged exposure to fumes. Wear natural rubber gloves, protective clothing and goggles when handling the concentrate to prepare dilutions. Wear a full-face mask of a type passed by the U. S. Department of Agriculture for DDVP (VAPONA Insecticide) protection. Keep all unprotected persons out of operational area.

In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted emulsion. It is recommended that only the amount of diluted emulsion be prepared from the concentrate that will be used immediately.

Mix the concentrate into water outdoors or in a well ventilated room. Wash off the outside of the bottle of concentrate in running water before storing away.

Do not contaminate lakes and streams.

Do not contaminate feed, water or foodstuffs.

Before applying in enclosed areas, remove animals from enclosures and reduce air movement as much as possible by closing doors, windows, and other openings. Do not treat animals or humans. Do not contaminate water, feed or foodstuffs. Ventilate before allowing animals to re-occupy treated premises. Repeat as necessary.

ANTIDOTE: If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY.**

If inhaled — Remove victim to fresh air. If on skin — Wash with large amounts of soap and water.

Do not re-use container. Destroy by rinsing thoroughly with water and alkaline detergent and crushing or perforating and burying in a safe place. Dispose of rinsings in a way as not to constitute a hazard or contaminate water supplies.

WARRANTY: Seller makes no warranty expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

U. S. D. A. Reg. No. 572-218