

Rockland 
CHEMICAL CO., INC.
Products for the Lawn, Garden, Farm, Home

PASSAIC AVENUE, WEST CALDWELL, N. J. / mailing address
P. O. BOX 204, CALDWELL, N. J. 07006
telephone (201) 575-1322 3 4

December 10, 1974

TO: Insecticide/Herbicide Branch
Environmental Protection Agency
Washington, D. C. 20460



RE: Additional insect control on label for EPA Reg. No. 572-210
PROFESSIONAL SUPER DUREBAN GRUBBY MUG KILLER

PROPOSED ADDITIONS:

The following is to be inserted on attached label, where indicated by arrow:

To control WHITE GRUBS (larvae of Japanese Beetle and European Chafer), use 2-4 lbs. per 1,000 sq. ft. Apply the granules uniformly over the area treated into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local agricultural extension service specialist for proper time to treat in your area.

RE Form 9-138 attached.



SUPER

PROFESSIONAL DURSBAN® CHINCH BUG KILLER

TREATS 25,000 SQ. FT.

ACCEPTED

MAY 30 1975

Under the Federal Insecticide,
Fungicide and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No.

For Professional Turf Maintenance Operators Only

A Dursban*-coated corn cob granular for control of Ants, Brown Dog Ticks, Chinch Bugs, Crickets, Curworms, Earwigs, Hyperodes (Turfgrass) Weevils, Sod Webworms (Lawn Moths) and Sowbugs (Pillbugs) on lawns and around buildings. Manufactured specifically for use in cyclone type spreaders.

DIRECTIONS FOR USE

Spread evenly at the rate of 1 pound per 1000 square feet. After application, water lawn using 30 gallons per 1000 square feet.

To control, ANTS, BROWN DOG TICKS, CRICKETS, CUTWORMS, EARWIGS, SOD WEBWORMS and SOWBUGS — Make first application when pests appear and re-treat as needed. For best results, turf should be moist at time of treatment. To control CHINCH BUGS — Consult your local state agricultural extension agent as to the exact timing.

NOTE: To reduce initial infestation or reinfestation of buildings by Earwigs, treat a band of ground 6 to 10 feet wide around and adjacent to the buildings where these pests are active.

CYCLONE SPREADER SETTING: # 3

To control HYPERODES WEEVILS — Apply as above except at twice the rate, or 2 pounds per 1000 sq. ft. Apply over suspected problem area in mid-April and again in mid-May.

HYPERODES WEEVIL SPREADER SETTING: #4½

These spreader settings are meant as a guide. They are in no sense a guarantee of exact application, inasmuch as age, wear of spreader, speed of operation and roughness of terrain can cause variations in rates applied.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not take internally. Avoid contact with skin, eyes or clothing. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before re-use. Avoid breathing of dust. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Keep away from food, feedstuffs and domestic water supplies.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Dispose of empty container and wastes by burning or burying in non-croplands away from domestic water supplies.

ACTIVE INGREDIENTS:

Chlorpyrifos [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate]	2.32%
Aromatic petroleum derivative solvent	1.26%
INERT INGREDIENTS:	96.42%

100.00%

E.P.A. Reg No. 572-219

*Dursban® Trademark of The Dow Chemical Co.
U.S. Pat. No. 3,244,586

E.P.A. Est. 572-N12

Net Weight 25 lbs.

Manufactured by

