



ROCKLAND

ONE QUART

# ROCKLAND FLOWABLE SEVIN\*

1  
2  
3

ACTIVE INGREDIENTS:  
Carbaryl (1-naphthyl N-methylcarbamate) 41.8%  
(Contains 4 lbs. Sevin per gallon)  
INERT INGREDIENTS: 58.2%  
\*Trade Mark Union Carbide Corp. for  
1-naphthyl N-methylcarbamate.

6-24-70  
572-221

## DIRECTIONS FOR USE

Before using shake or stir well. Pour recommended amount of this material into nearly filled spray tank, or pre-mix in bucket half-filled with water and pour mixture into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Use a screen not finer than 50 mesh in entire system. Do not combine with emulsifiable liquids. Do not use with lime, Bordeaux, lime-sulfur, casein-lime spreaders or other alkaline materials. Some phytotoxicity may occur on tender foliage when prolonged misty rain or high humidity follows spraying. Does not control spider mites, but is compatible with all common miticides.

**ORNAMENTALS:** To control apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale crawlers, tent caterpillars, thrips (exposed), willow leaf beetles on Rose, Carnation, Gladiolus, Zinnia, Chrysanthemum, Lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines (do not use on Boston Ivy). Apply 1 qt. per 100 gal. water (2 teaspoonfuls per gal.) when insects or their damage appear. Repeat weekly or as needed.

**LAWN INSECTS:** To control ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, sod webworms (lawn moths), millipeds. Apply 4 oz. in 15 to 20 gal. water for each 500 sq. ft. of established lawn area. Use garden hose sprayers or pressure type equipment and apply full-water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat in 2-3 weeks if necessary.

**POULTRY PESTS:** To control chicken mite, fleas, lice and northern fowl mite on Chickens, Ducks, Game Birds, Geese, Pigeons, Turkeys, direct spray mist on birds. Do not apply within 7 days of slaughter. Do not repeat application within 4 weeks. Do not contaminate feed or drinking water. This is a supplement to spraying roosts and buildings for control of chicken mites and fleas.

**MISTING WITH ELECTRIC FOG MACHINE:** Mix 10 fl. oz. in 1 gal. spray. Use 1 1/2 gal. per 1,000 hens in cages, on litter or on slatted floor. Repeat in 4 weeks if necessary.

**SPRAY WITH KNAPSACK OR CYLINDER TYPE COMPRESSED AIR SPRAYERS:** Mix 6 fl. oz. in 5 gal. spray. Use 1 gal. per 100 hens in cages on litter or on slatted floor. Repeat in 4 weeks if necessary.

**SPRAY ROOSTS AND BUILDINGS** with conventional power spray or knapsack equipment. **FOR BEDBUGS, CHICKEN MITES, FLEAS,** use 1 qt. per 25 gal. water. **FOR FOWL TICKS,** use 1 gal. per 25 gal. water. Spray 1/2 gal. per 1,000 sq. ft. of wall, bedding, litter or roost surfaces. Force spray into cracks and crevices. Repeat as needed. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water. Do not house birds in treated houses within 7 days of slaughter.

**Caution:** Allow 7 days between application to poultry and game birds or poultry and game bird premise treatments and bird slaughter. Avoid contamination of nests, eggs and feeding and watering troughs. Ventilate while spraying.

**CAUTION:** Harmful if swallowed or inhaled. Avoid prolonged or repeated breathing of spray. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Take shower or bath after use. Wear regular long sleeved work clothing. Change to clean clothing daily. **NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of food, feed, feeding troughs and water receptacles.

This product is highly toxic to bees exposed to direct treatment or residues in crops. For protection of honeybees avoid unnecessary use during periods when honeybees are visiting the crop. Avoid spray drift to adjoining food and forage crops or crops in bloom. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured for:

ROCKLAND CHEMICAL CO., Inc.  
PASSAIC AVENUE WEST CALDWELL, N. J.

U.S.D.A. Reg. No. 572-221



ROCKLAND

ONE QUART

# ROCKLAND FLOWABLE SEVIN\*

1  
2  
3

ACTIVE INGREDIENTS:  
Carbaryl (naphthyl N-methyl carbamate) 41.8%  
(Contains 4 lbs. Sevin per gallon)  
INERT INGREDIENTS: 58.2%  
\*Trade Mark Union Carbide Corp. for  
1-naphthyl N-methyl carbamate.

6-24-70  
572-221

### DIRECTIONS FOR USE

Before using shake or stir well. Pour recommended amount of this material into nearly filled spray tank, or pre-mix in bucket half-filled with water and pour mixture into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Use a screen not finer than 50 mesh in entire system. Do not combine with emulsifiable liquids. Do not use with lime, Bordeaux, lime-sulfur, casein-lime spreaders or other alkaline materials. Some phytotoxicity may occur on tender foliage when prolonged misty rain or high humidity follows spraying. Does not control spider mites, but is compatible with all common miticides.

**ORNAMENTALS:** To control apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bog, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale crawlers, tent caterpillars, thrips (exposed), willow leaf beetles on Rose, Carnation, Gladiolus, Zinnia, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, Birch and pines (do NOT use on Boston Ivy). Apply 1 qt. per 100 gal. water (2 teaspoonfuls per gal.) when insects or their damage appear. Repeat weekly or as needed.

**LAWN INSECTS:** To control ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, sod webworms (law moths), millipeds, Apply 4 oz. in 15 to 20 gal. water for each 500 sq. ft. of established lawn area. Use garden hose sprayers or pressure type equipment and apply full-water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat in 2-3 weeks if necessary.

**POULTRY PESTS:** To control chicken mite, fleas, lice and northern fowl mite on Chickens, Ducks, Game Birds, Geese, Pigeons, Turkeys, direct spray mist on birds. Do not apply within 7 days of slaughter. Do not repeat application within 4 weeks. Do not contaminate feed or drinking water. This is a supplement to spraying roosts and buildings for control of chicken mites and fleas.

**MISTING WITH ELECTRIC FOG MACHINE:** Mix 10 fl. oz. in 1 gal. spray. Use 1/2 gal. per 1,000 hens in cages, on litter or on slatted floor. Repeat in 4 weeks if necessary.

**SPRAY WITH KNAPSACK OR CYLINDER TYPE COMPRESSED AIR SPRAYERS:** Mix 6 fl. oz. in 5 gal. spray. Use 1 gal. per 100 hens in cages on litter or on slatted floor. Repeat in 4 weeks if necessary.

**SPRAY ROOSTS AND BUILDINGS** with conventional power spray or knapsack equipment. **FOR BEDBUGS, CHICKEN MITES, FLEAS,** use 1 qt. per 25 gal. water. **FOR FOWL TICKS,** use 1 gal. per 15 gal. water. Spray 1/2 gal. per 1,000 sq. ft. of wall, bedding, litter or roost surfaces. Force spray into cracks and crevices. Repeat as needed. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water. Do not house birds in treated houses within 7 days of slaughter.

**Caution:** Allow 7 days between application to poultry and game birds or poultry and game bird premise treatments and bird slaughter. Avoid contamination of nests, eggs and feeding and watering troughs. Ventilate while spraying.

**CAUTION:** Harmful if swallowed or inhaled. Avoid prolonged or repeated breathing of spray. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Take shower or bath after use. Wear regular long sleeved work clothing. Change to clean clothing daily. **NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of food, feed, feeding troughs and water receptacles.

This product is highly toxic to bees exposed to direct treatment or residues in crops. For protection of honeybees avoid unnecessary use during periods when honeybees are visiting the crop. Avoid spray drift to adjoining food and forage crops or crops in bloom. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured for:

ROCKLAND CHEMICAL CO., Inc.

PASSAIC AVENUE

WEST CALDWELL, N. J.

U.S.D.A. Reg. No. 572-221



ROCKLAND

ONE QUART

# ROCKLAND FLOWABLE SEVIN\*

1  
2  
3

ACTIVE INGREDIENTS:  
Cyfluthrin 4.0%  
Inert Ingredients 96.0%  
Net Weight 1.05 Quarts

6-24-70  
572-221

### DIRECTIONS FOR USE

Before using shake or stir well. Pour recommended amount of this material into nearly filled spray tank or pre-mix in bucket half-filled with water and pour mixture into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Use a screen not finer than 50 mesh in entire system. Do not combine with emulsifiable liquids. Do not use with lime, Bordeaux, lime-sulfur, casein-lime spreaders or other alkaline materials. Some phytotoxicity may occur on tender foliage when prolonged misty rain or high humidity follows spraying. Does not control spider mites, but is compatible with all common miticides.

**ORNAMENTALS:** To control apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale crawlers, tent caterpillars, thrips (exposed), willow leaf beetles on Rose, Carnation, Gladiolus, Zinnia, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines (do NOT use on Boston Ivy). Apply 1 qt. per 100 gal. water (2 teaspoonfuls per gal.) when insects or their damage appear. Repeat weekly or as needed.

**LAWN INSECTS:** To control ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, sod webworms (lawn moths), millipedes. Apply 4 oz. in 15 to 20 gal. water for each 500 sq. ft. of established lawn area. Use garden hose sprayers or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat in 2-3 weeks if necessary.

**POULTRY PESTS:** To control chicken mite, fleas, lice and northern fowl mite on Chickens, Ducks, Game Birds, Geese, Pigeons, Turkeys, direct spray mist on birds. Do not apply within 7 days of slaughter. Do not repeat application within 4 weeks. Do not contaminate feed or drinking water. This is a supplement to spraying roosts and buildings for control of chicken mites and fleas.

**MISTING WITH ELECTRIC FOG MACHINE:** Mix 10 fl. oz. in 1 gal. spray. Use 1 1/2 gal. per 1,000 hens in cages on litter or on slatted floor. Repeat in 4 weeks if necessary.

**SPRAY WITH KNAPSACK OR CYLINDER TYPE COMPRESSED AIR SPRAYERS:** Mix 6 fl. oz. in 1 gal. water. Use 1 gal. per 100 hens in cages on litter or on slatted floor. Repeat in 4 weeks if necessary.

**SPRAY ROOSTS AND BUILDINGS** with concentration shown spray or mist as needed. **FOR BEDBUGS, CHICKEN MITES, FLEAS,** use 1 qt. per 25 gal. water. **FOR FOWL TICKS,** use 1 qt. per 5 gal. water. Spray on litter or 1,000 hens in cages on bedding, litter or roost surfaces. Force spray into cracks and crevices. Repeat as needed. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water. Do not house birds in treated houses within 7 days of slaughter.

**Caution:** Allow 7 days between application to poultry and game birds or to any and game bird premise treatments and bird slaughter. Avoid contamination of feed, eggs and feeding and watering troughs. Ventilate when spraying.

**CAUTION:** Harmful if swallowed or inhaled. Avoid prolonged or repeated breathing of spray. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Take shower or bath after use. Wear regular long sleeved work clothing. Change to clean clothing daily. **NOTE TO PHARMACIANS:** Carbarvyl is a moderate, reversible, non-competitive inhibitor. Atropine is antidotal. Avoid contamination of food, feed, feeding troughs and water receptacles.

This product is highly toxic to bees exposed to direct treatment or residues on crops. For protective purposes, bees avoid unnecessary use during periods when bees are visiting the crop. Avoid spraying when bees are visiting food and storage areas on crops. For more information, may be obtained from your local Agricultural Extension Service.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured for:

ROCKLAND CHEMICAL CO., Inc.

PASSAIC AVENUE

WEST CALDWELL, N. J.