

[572-224]

**SPOT SPRAYS:** For localized housefly control—Apply a 1% spray with a knapsack or similar type sprayer. Apply spray to areas where flies congregate such as doorways and around window sills. Repeat as necessary.

**MAGGOT SPRAYS:** For control of housefly maggots—Apply a 1% spray to poultry droppings, garbage dumps, manure piles, etc. Repeat applications as additional manure or garbage is added.

**GENERAL OUTSIDE USE:** For control of houseflies around homes and recreation areas, garbage cans, animal quarters, food processing plants, warehouses, loading docks and garbage dumps. Apply a 1% spray to exposed areas. Repeat applications as necessary.

**CAUTION:** May cause irritation of eyes, nose, throat and skin. May be harmful if inhaled or swallowed, or absorbed through the skin. Vapor harmful. Avoid breathing vapor or spray mist. Use only with adequate ventilation. Avoid contact with eyes and skin. Wash thoroughly after handling.

In case of contact, flush with plenty of water: for eyes, get medical attention.

**NOTE TO PHYSICIAN:** This product upon repeated, prolonged or careless use, may cause Cholinesterase Inhibition. Antidote is ATROPINE.

Do not contaminate or apply onto feed and foodstuffs, drinking fountains, litter and feed troughs. Do not use in milk processing rooms including milk houses and milk storage rooms. Do not use in homes. Keep out of reach of domestic animals and away from food stuffs.

Birds and other wildlife feeding on treated areas may be killed.

Apply this product only as specified on this label. Do not re-use empty container. Destroy container by perforating or crushing and burying in a safe place.

U.S.D.A. Reg. No. 572-224  
I.B.A. Stock No. 4938

NET CONTENTS ONE GALLON



# RESIDUAL FLY SPRAY 8

An insecticide used for controlling house flies and maggots in Dairy Barns, Poultry Houses, Hog Pens and certain other farm buildings.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

See Back Panel for Additional Cautions

**ACTIVE INGREDIENT:**  
Dimethoate (O,O-Dimethyl S-(N-Methylcarbamoyl)methyl) Phosphorodithioate 73.4%  
(Equivalent to 2.1 lbs. per gallon)  
**INERT INGREDIENTS:** 75.6%

MANUFACTURED FOR  
**I. B. A.**  
MILLBURY, MASS.

IBA RESIDUAL FLY SPRAY 8 will control flies up to 1000 feet. IBA RESIDUAL FLY SPRAY should be applied as follows:  
How to prepare a 1% RESIDUAL FLY SPRAY:

AMOUNT OF SPRAY NEEDED	AMOUNT OF RESIDUAL FLY SPRAY 8 NEEDED
1 Pint	4 Teaspoons
2 Quarts	1/2 Cup
6 Quarts	1/2 Pint
7 Gallons	1 Pint
6 Gallons	1 Quart
12 Gallons	2 Quarts
25 Gallons	1 Gallon

IBA RESIDUAL FLY SPRAY 8 is a clear liquid. When mixed with water, it will disperse readily and will remain stable with a minimum amount of agitation.

**DIRECTIONS FOR USE:** IBA RESIDUAL FLY SPRAY 8 should be applied by a knapsack or power sprayer. As a wall spray, prepare a 1% solution of IBA RESIDUAL FLY SPRAY 8 and spray ceilings, walls and floors. One gallon of a 1% RESIDUAL FLY SPRAY 8 will cover 500 to 1000 sq. ft.

As a spot spray prepare a 1% spray of RESIDUAL FLY SPRAY 8, apply with a knapsack sprayer and apply to areas where flies congregate.

**RESIDUAL WALL SPRAYS:** For control of houseflies and resistant strains in Dairy Barns, Calf Barns, Poultry Houses and other farm buildings.

Apply a 1% spray. Thoroughly wet all fly resting surfaces to the point of run-off. Apply spray to ceilings, walls and floors. One gallon of a 1% spray will cover 500 to 1000 sq. ft. of surface. Repeat application when necessary.

**REMOVE DAIRY ANIMALS, CALVES, AND POULTRY FROM BUILDINGS PRIOR TO SPRAYING.**

ACCEPTED  
7-16-10  
572-224

72-224

or localized housefly control--Apply a 1% sack or similar type sprayer. Apply spray to congregate such as doorways and around it as necessary.

S: For control of housefly maggots--Apply poultry droppings, garbage dumps, manure piles, etc. Repeat applications as or garbage is added.

DE USE: For control of houseflies around lion areas, garbage cans, animal quarters, plants, warehouses, loading docks and gar- ly a 1% spray to exposed areas. Repeat essary.

ause irritation of eyes, nose, throat and nful if inhaled or swallowed, or absorbed vapor harmful. Avoid breathing vapor or y with adequate ventilation. Avoid con- skin. Wash thoroughly after handling.

flush with plenty of water; for eyes, get

IAN: This product upon repeated, pro- use, may cause Cholinesterase Inhibition. INE.

or apply onto feed and foodstuffs, drink- r and feed troughs. Do not use in milk including milk houses and milk storage in homes. Keep out of reach of domestic om food stuffs.

idlife feeding on treated areas may be

only as specified on this label. y container. Destroy container by perfor- burying in a safe place.

.D.A. Reg. N- 572 224  
IBA Stock No. 4938

NET CONTENTS ONE GALLON



# RESIDUAL FLY SPRAY 8

An insecticide used for controlling house flies and maggots in Dairy Barns, Poultry Houses, Hog Pens and certain other farm buildings.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
See Back Panel for Additional Cautions

ACTIVE INGREDIENT:  
Dimethoate (O,O-Dimethyl S-(N-Methylcarbamoylmethyl) Phosphorodithioate) 73.4%  
(Equivalent to 2.1 lbs. per gallon)  
INERT INGREDIENTS: 75.6%

MANUFACTURED FOR  
**I. B. A.**  
MILLBURY, MASS.

IBA RESIDUAL FLY SPRAY 8 will control flies up to 8 weeks. IBA RESIDUAL FLY SPRAY should be applied as a 1% spray. How to prepare a 1% RESIDUAL FLY SPRAY 8:

AMOUNT OF SPRAY NEEDED	AMOUNT OF RESIDUAL FLY SPRAY 8 NEEDED
1 Pint	4 Teaspoons
2 Quarts	1/2 Cup
6 Quarts	1/2 Pint
3 Gallons	1 Pint
6 Gallons	1 Quart
12 Gallons	2 Quarts
25 Gallons	1 Gallon

IBA RESIDUAL FLY SPRAY 8 is a clear liquid. When added to water, it will disperse readily and will remain physically stable with a minimum amount of agitation.

DIRECTIONS FOR USE: IBA RESIDUAL FLY SPRAY 8 can be applied by a knapsack or power sprayer. As a residual wall spray prepare a 1% solution of IBA RESIDUAL FLY SPRAY 8 and spray ceilings, walls and floors. One gallon of a 1% RESIDUAL FLY SPRAY 8 will cover 500 to 1000 square feet.

As a spot spray prepare a 1% spray of RESIDUAL FLY SPRAY 8, apply with a knapsack sprayer and apply wherever flies congregate.

RESIDUAL WALL SPRAYS: For control of houseflies, including resistant strains in Dairy Barns, Calf Barns, Hog Pens, Poultry Houses and other farm buildings.

Apply a 1% spray. Thoroughly wet all fly resting areas to the point of run-off. Apply spray to ceilings, walls and floors. One gallon of a 1% spray will cover 500 to 1000 square feet of surface. Repeat application when necessary.

REMOVE DAIRY ANIMALS, CALVES, AND POULTRY FROM BUILDINGS PRIOR TO SPRAYING.

ACCEPTED  
7-16-70  
572-224

[572-224]

**SPOT SPRAYS:** For localized housefly control—Apply a 1% spray with a knapsack or similar type sprayer. Apply spray to areas where flies congregate such as doorways and around window sills. Repeat as necessary.

**MAGGOT SPRAYS:** For control of housefly maggots—Apply a 1% spray to poultry droppings, garbage dumps, manure piles ~~and other areas where they are found~~. Repeat applications as additional manure or garbage is added.

**GENERAL OUTSIDE USE:** For control of houseflies around homes and recreation areas, garbage cans, animal quarters, food processing plants, warehouses, loading docks and garbage dumps. Apply a 1% spray to exposed areas. Repeat applications as necessary.

**CAUTION:** May cause irritation of eyes, nose, throat and skin. May be harmful if inhaled or swallowed, or absorbed through the skin. Vapor harmful. Avoid breathing vapor or spray mist. Use only with adequate ventilation. Avoid contact with eyes and skin. Wash thoroughly after handling.

In case of contact, flush with plenty of water; for eyes, get medical attention.

**NOTE TO PHYSICIAN:** This product upon repeated, prolonged or careless use, may cause Cholinesterase Inhibition. Antidote is ATROPINE.

Do not contaminate or apply onto feed and foodstuffs, drinking fountains, litter and feed troughs. Do not use in milk processing rooms including milk houses and milk storage rooms. ~~Do not use in homes. Keep out of reach of domestic animals and away from food stuffs.~~

~~Do not use in homes. Keep out of reach of domestic animals and away from food stuffs.~~


Birds and other wildlife feeding on treated areas may be killed.

Apply this product only as specified on this label.

Do not re-use empty container. Destroy container by perforating or crushing and burying in a safe place.

U.S.D.A. Reg. No. 572-224

BA Stock No. 4938




**RESIN**

An insecticide  
Barns, Poultry

**CAUTION**

ACTIVE INGREDIENTS  
Dimethoate  
Phosphorus  
(Equivalent)  
INERT INGREDIENTS

CONTENTS ONE GALLON



# RESIDUAL FLY SPRAY 8

For controlling house flies and maggots in Dairy Barns, Hog Pens and certain other farm buildings.

**KEEP OUT OF REACH OF CHILDREN.**

See Back Panel for Additional Cautions

IBA: .....  
 (N-Dimethyl S-(N-Methylcarbamoylmethyl) 23.4%  
 (e)  
 (2.1 lbs. per gallon)  
 IBS: .....75.6%

MANUFACTURED FOR

**I. B. A.**  
MILLBURY, MASS.

IBA RESIDUAL FLY SPRAY 8 will control flies up to 8 weeks. IBA RESIDUAL FLY SPRAY should be applied as a 1% spray. How to prepare a 1% RESIDUAL FLY SPRAY 8:

AMOUNT OF SPRAY NEEDED	AMOUNT OF RESIDUAL FLY SPRAY 8 NEEDED
1 Pint	4 Teaspoons
2 Quarts	1/3 Cup
6 Quarts	1/2 Pint
3 Gallons	1 Pint
6 Gallons	1 Quart
12 Gallons	2 Quarts
25 Gallons	1 Gallon

IBA RESIDUAL FLY SPRAY 8 is a clear liquid. When added to water, it will disperse readily and will remain physically stable with a minimum amount of agitation.

**DIRECTIONS FOR USE:** IBA RESIDUAL FLY SPRAY 8 can be applied by a knapsack or power sprayer. As a residual wall spray, prepare a 1% solution of IBA RESIDUAL FLY SPRAY 8 and spray ceilings, walls and floors. One gallon of a 1% RESIDUAL FLY SPRAY 8 will cover 500 to 1000 square feet.

As a spot spray prepare a 1% spray of RESIDUAL FLY SPRAY 8, apply with a knapsack sprayer and apply wherever flies congregate.

**RESIDUAL WALL SPRAYS:** For control of houseflies, including resistant strains in Dairy Barns, Calf Barns, Hog Pens, Poultry Houses and other farm buildings.

Apply a 1% spray. Thoroughly wet all fly resting areas to the point of run-off. Apply spray to ceilings, walls and floors. One gallon of a 1% spray will cover 500 to 1000 square feet of surface. Repeat application when necessary.

**REMOVE DAIRY ANIMALS, CALVES, AND POULTRY FROM BUILDINGS PRIOR TO SPRAYING.**